

ON THE RATE OF CONVERGENCE OF FINITE-DIFFERENCE APPROXIMATIONS FOR NORMALIZED BELLMAN EQUATIONS WITH LIPSCHITZ COEFFICIENTS

ISTVÁN GYÖNGY, DAVID ŠIŠKA

ABSTRACT. A class of stochastic optimal control problems containing optimal stopping of controlled diffusion process is considered. The numerical solutions of the corresponding normalized Bellman equations are investigated. Methods of [1] are adapted. The rate of convergence of appropriate finite difference difference schemes is estimated.

1. INTRODUCTION

Stochastic optimal control and optimal stopping problems have many applications in mathematical finance, portfolio optimization, economics and statistics (sequential analysis). Optimal stopping problems can be in some cases solved analytically [2]. With most problems, one must resort to numerical approximations of the solutions. One approach is to use controlled Markov chains as an approximation to the controlled diffusion process, see e.g. [3]. A thorough account of this approach is available in [4].

We are interested in the rate of convergence of finite difference approximations to the payoff function of optimal control problems when the reward and discounting functions may be unbounded in the control parameter. This allows us to treat numerically the optimal stopping of controlled diffusion processes by randomized stopping, i.e. by transforming the optimal stopping into a control problem, see [5]. This leads us to approximating a normalized degenerate Bellman equation.

Until quite recently, there were no results on the rate of convergence of finite difference schemes for degenerate Bellman equations. A major breakthrough is achieved in Krylov [6] for Bellman equations with constant coefficients, followed by rate of convergence estimates for Bellman equations with variable coefficients in [7] and [8]. The estimate from [8] is improved in [9] and [10]. Finally, Krylov [1] (published in [11]) establishes the rate of convergence $\tau^{1/4} + h^{1/2}$ of finite difference schemes to degenerate Bellman equations with Lipschitz coefficients, where τ and h are the mesh sizes in time and space respectively.

In the present paper we extend this estimate to a class of normalized degenerate Bellman equations which contains those for optimal stopping of controlled diffusion processes with variable coefficients. Adapting ideas and techniques of [1] we obtain the rate of convergence $\tau^{1/4} + h^{1/2}$, as in [1]. The main ingredient of the proof is a gradient estimate for the solution to the discrete normalized Bellman PDE. This is an extension of the gradient estimate from [1] to our case. After the first version of this article was sent to

arxiv, we received [12] where an essentially more general gradient estimate is proved. This opens the way to proving the same rate of convergence result for normalized Bellman PDEs in more general setting then we present below.

Rate of converge results for optimal stopping are proved for general consistent approximation schemes in [13]. However, the rate $\tau^{1/4} + h^{1/2}$ is obtained only when the diffusion coefficients are independent of the time and space variables. For further results on numerical approximations for Bellman equations we refer to [14], [15] and [16].

The paper is organized as follows. The main result is formulated in the next section. In section 3 the existence and uniqueness of the solution to finite difference schemes is proved together with a comparison result. The main technical result, the gradient estimate of solutions to finite difference schemes, is obtained in section 4. Some useful analytic properties of payoff functions are presented in section 5. Finally, the main result is proved in section 6.

2. THE MAIN RESULT

Fix $T \in [0, \infty)$. Let $(\Omega, \mathcal{F}, \mathbb{P}, (\mathcal{F}_t)_{t \geq 0})$ be a probability space with a right-continuous filtration, such that $\mathcal{F}_0$ contains all $\mathbb{P}$ null sets. Let $(w_t, \mathcal{F}_t)$ be a d' dimensional Wiener martingale. Let A be a separable metric space. For every $t \in [0, T]$, $x \in \mathbb{R}^d$ and $\alpha \in A$ we are given a $d \times d'$ dimensional matrix $\sigma^\alpha(t, x)$, a d dimensional vector $\beta^\alpha(t, x)$ and real numbers $c^\alpha(t, x)$, $f^\alpha(t, x)$ and $g(x)$.

Assumption 2.1. σ, β, c, f are Borel functions of (α, t, x) . The function g is continuous in x . There exist an increasing sequence of subsets A_n of A , and positive real constants K, K_n , and m, m_n , such that $\bigcup_{n \in \mathbb{N}} A_n = A$ and for each $n \in \mathbb{N}$, $\alpha \in A_n$,

$$\begin{aligned} |\sigma^\alpha(t, x) - \sigma^\alpha(t, y)| + |\beta^\alpha(t, x) - \beta^\alpha(t, y)| &\leq K_n |x - y|, \\ |\sigma^\alpha(t, x)| + |\beta^\alpha(t, x)| &\leq K_n (1 + |x|), \\ |c^\alpha(t, x)| + |f^\alpha(t, x)| &\leq K_n (1 + |x|)^{m_n}, \quad |g(x)| \leq K (1 + |x|)^m \end{aligned} \quad (2.1)$$

for all $x \in \mathbb{R}^d$ and $t \in [0, T]$.

We say that $\alpha \in \mathfrak{A}_n$ if $\alpha = (\alpha_t)_{t \geq 0}$ is a progressively measurable process with values in A_n . Let $\mathfrak{A} = \bigcup_{n \in \mathbb{N}} \mathfrak{A}_n$. Then under Assumption 2.1 it is well known that for each $s \in [0, T]$, $x \in \mathbb{R}^d$ and $\alpha \in \mathfrak{A}$ there is a unique solution $\{x_t : t \in [0, T - s]\}$ of

$$x_t = x + \int_0^t \sigma^{\alpha_u}(s + u, x_u) dw_u + \int_0^t \beta^{\alpha_u}(s + u, x_u) du, \quad (2.2)$$

denoted by $x_t^{\alpha, s, x}$. For $s \in [0, T]$ we use the notation $\mathfrak{T}(T - s)$ for the set of stopping times $\tau \leq T - s$. Define the payoff function to the optimal stopping and control problem as

$$w(s, x) = \sup_{\alpha \in \mathfrak{A}} \sup_{\tau \in \mathfrak{T}(T-s)} v^{\alpha, \tau}(s, x), \quad (2.3)$$

where

$$v^{\alpha, \tau}(s, x) = \mathbb{E}_{s, x}^\alpha \left[\int_0^\tau f^{\alpha_t}(s + t, x_t) e^{-\varphi_t} dt + g(x_\tau) e^{-\varphi_\tau} \right],$$

$$\varphi_t = \int_0^t c^{\alpha_r}(s+r, x_r) dr,$$

and $\mathbb{E}_{s,x}^\alpha$ means expectation of the expression behind it, with $x_t^{\alpha,s,x}$ in place of x_t everywhere. It is worth noticing that for

$$w_n(s, x) := \sup_{\alpha \in \mathfrak{A}_n} \sup_{\tau \in \mathfrak{T}(T-s)} v^{\alpha, \tau}(s, x)$$

we have $w_n(s, x) \uparrow w(s, x)$ as $n \rightarrow \infty$. By Theorem 3.1.8 in [5], $w_n(s, x)$ is bounded from above and below. Hence $w(s, x)$ is bounded from below. However it can be equal to $+\infty$.

Let $\mathfrak{R}_n$ contain all progressively measurable, locally integrable processes $r = (r_t)_{(t \geq 0)}$ taking values in $[0, n]$ such that $\int_0^\infty r_t dt = \infty$. Let $\mathfrak{R} = \bigcup_{n \in \mathbb{N}} \mathfrak{R}_n$.

We will state without proof a result about randomized stopping. The proof (taking into account the possibility that g is a function of (t, x) , continuous in (t, x)) can be found in [17]. If $A = A_n$, $K = K_n$, $m = m_n$ for $n \geq 1$ then the following theorem is known from [5] (see Exercise 3.4.12, via Lemma 3.4.3(b) and Lemma 3.4.5(c)).

Theorem 2.2. *Let Assumption 2.1 hold. Then for all $(s, x) \in [0, T] \times \mathbb{R}^d$, either both $w(s, x)$ and*

$$\begin{aligned} \sup_{\alpha \in \mathfrak{A}} \sup_{r \in \mathfrak{R}} \mathbb{E}_{s,x}^\alpha \left\{ \int_0^{T-s} [f^{\alpha_t}(s+t, x_t) e^{-\varphi_t} \right. \\ \left. + r_t g(x_t) e^{-\varphi_t}] e^{-\int_0^t r_u du} dt + g(x_{T-s}) e^{-\varphi_{T-s} - \int_0^{T-s} r_u du} \right\} \end{aligned} \quad (2.4)$$

are finite and equal, or they are both infinite.

This theorem is the main tool which allows us to treat the problem of optimal stopping of a controlled process as just an optimal control problem. The aim of this paper is to find the rate of convergence for numerical approximations to the following payoff function:

$$v(s, x) = \sup_{\alpha \in \mathfrak{A}} v^\alpha(s, x), \quad (2.5)$$

$$v^\alpha(s, x) = \mathbb{E}_{s,x}^\alpha \left[\int_0^{T-s} f^{\alpha_t}(s+t, x_t) e^{-\varphi_t} dt + g(x_{T-s}) e^{-\varphi_{T-s}} \right], \quad (2.6)$$

where x_t is the solution to (2.2). Notice that (2.4) can be written in this form such that Assumption 2.1 is satisfied.

The immediate aim now is to find an approximation scheme for the payoff function. From [5] we know that under some assumptions (stricter than Assumptions 2.1) the payoff function (2.5) satisfies the normalized Bellman equation

$$\begin{aligned} \sup_{\alpha \in A} m^\alpha(u_t + L^\alpha u + f^\alpha) &= 0 \quad \text{on } [0, T] \times \mathbb{R}^d \\ u(T, x) &= g(x) \quad \text{for } x \in \mathbb{R}^d, \end{aligned} \quad (2.7)$$

where

$$L^\alpha u = \sum_{i,j} a_{i,j}^\alpha u_{x_i, x_j} + \sum_i \beta_i^\alpha u_{x_i} - c^\alpha u,$$

$a_{ij}^\alpha = 1/2(\sigma^\alpha \sigma^{\alpha*})_{ij}$ and m^α is a non-negative function of the control parameter called the normalizing factor. A crucial property of m^α is that $m^\alpha \sigma_{ij}^\alpha$, $m^\alpha b_i^\alpha$, $m^\alpha c^\alpha$ and $m^\alpha f^\alpha$ are bounded as functions α . Hence it is natural to approximate v by approximating the solution of the normalized Bellman equation. In this paper we show that the finite difference approximation to the solution of the normalized Bellman PDE converges to v . Moreover we find that the rate of convergence is the same as in [1]. It is worth noticing that to show this we don't need any result about the solvability of the Bellman equation. We now describe the approximation scheme. From now on let $K \geq 1$ be a fixed constant.

Assumption 2.3. There exist a natural number d_1 , vectors $\ell_k \in \mathbb{R}^d$ and functions

$$\sigma_k^\alpha : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}, \quad b_k^\alpha : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}, \quad \forall k = \pm 1, \dots, d_1$$

such that $\ell_k = \ell_{-k}$, $|\ell_k| \leq K$, $\sigma_k^\alpha = \sigma_{-k}^\alpha$, $b_k^\alpha \geq 0$ for $k = \pm 1, \dots, d_1$ and

$$\sigma_{ij}^\alpha(t, x) = \sigma_k^\alpha(t, x) \ell_k^i \ell_k^j, \quad \beta_i^\alpha(t, x) = b_k^\alpha(t, x) \ell_k^i \quad (2.8)$$

for any $\alpha \in A$ and $(t, x) \in [0, T] \times \mathbb{R}^d$.

This might appear to be a restrictive assumption. Actually, all operators L^α which admit monotone finite difference approximations satisfy this. See [18]. Fix $\tau > 0$, $h > 0$ and define the grid

$$\begin{aligned} \bar{\mathcal{M}}_T := \{(t, x) \in [0, T] \times \mathbb{R}^d : (t, x) = ((j\tau) \wedge T, h(i_1 \ell_1 + \dots + i_{d_1} \ell_{d_1})), \\ j \in \{0\} \cup \mathbb{N}, i_k \in \mathbb{Z}, k = \pm 1, \dots, \pm d_1\}. \end{aligned}$$

Let $\tau_T(t) := \tau$ for $t \leq T - \tau$ and $\tau_T(t) := T - t$ for $t > T - \tau$. So $t + \tau_T(t) = (t + \tau) \wedge T$. Let Q be a non-empty subset of

$$\mathcal{M}_T := \bar{\mathcal{M}}_T \cap ([0, T] \times \mathbb{R}^d).$$

Let $T_\tau u(t, x) = u(t + \tau_T(t), x)$, $T_{h, \ell_k} u(t, x) = u(t, x + h_k \ell_k)$,

$$\delta_\tau u(t, x) = \frac{u(t + \tau_T(t), x) - u(t, x)}{\tau_T(t)},$$

$$\delta_{h_k, \ell_k} u(t, x) = \frac{u(t, x + h_k \ell_k) - u(t, x)}{h_k} \quad \text{and} \quad \Delta_{h_k, \ell_k} u = -\delta_{h_k, \ell_k} \delta_{h_k, -\ell_k} u.$$

Let $\delta_{0, \ell} := 0$. Let $a_k^\alpha := (1/2)(\sigma_k^\alpha)^2$. Consider the following finite difference problem:

$$\sup_{\alpha \in A} m^\alpha (\delta_\tau u + L_h^\alpha u + f^\alpha) = 0 \quad \text{on} \quad Q, \quad (2.9)$$

$$u = g \quad \text{on} \quad \bar{\mathcal{M}}_T \setminus Q, \quad (2.10)$$

where

$$L_h^\alpha u = \sum_k a_k^\alpha \Delta_{h_k, \ell_k} u + \sum_k b_k^\alpha \delta_{h_k, \ell_k} u - c^\alpha u$$

and m^α is a positive function of $\alpha \in A$ such that the following conditions hold.

Assumption 2.4. Let $0 \leq \lambda < \infty$. Functions $\sigma_k^\alpha, b_k^\alpha, f^\alpha, c^\alpha \geq \lambda, g$ are Borel in t and continuous in $\alpha \in A$ for each $k = \pm 1, \dots, d_1$. For any $t \in [0, T]$, $x, y \in \mathbb{R}^d$ and $\alpha \in A$:

$$\begin{aligned} & |\sigma_k^\alpha(t, x) - \sigma_k^\alpha(t, y)| + |b_k^\alpha(t, x) - b_k^\alpha(t, y)| + |c^\alpha(t, x) - c^\alpha(t, y)| \\ & + m^\alpha |f^\alpha(t, x) - f^\alpha(t, y)| + |g(x) - g(y)| \leq K|x - y|, \quad (2.11) \\ & |\sigma_k^\alpha| + b_k^\alpha + m^\alpha |f^\alpha| + m^\alpha c^\alpha + |g| \leq K \end{aligned}$$

and

$$\frac{1}{K} \leq m^\alpha [1 + c^\alpha(t, x)]. \quad (2.12)$$

Notice that if Assumptions 2.3 and 2.4 are satisfied then (2.1) is also satisfied. On top of Assumptions 2.1, 2.3 and 2.4, the following is required to obtain the rate of convergence.

Assumption 2.5. For all $\alpha \in A, x \in \mathbb{R}^d$ and $t, s \in [0, T]$

$$\begin{aligned} & |\sigma^\alpha(s, x) - \sigma^\alpha(t, x)| + |b^\alpha(s, x) - b^\alpha(t, x)| \\ & + |c^\alpha(s, x) - c^\alpha(t, x)| \leq K|t - s|^{1/2}, \\ & m^\alpha |f^\alpha(s, x) - f^\alpha(t, x)| \leq K|t - s|^{1/2}. \end{aligned}$$

The following theorem is the main result of the paper.

Theorem 2.6. *Let Assumptions 2.1, 2.3, 2.4 and 2.5 be satisfied. Let v be the payoff function (2.5). Then (2.9)-(2.10) with $Q = \mathcal{M}_T$ has a unique bounded solution $v_{\tau, h}$ and*

$$|v - v_{\tau, h}| \leq N_T(\tau^{1/4} + h^{1/2}) \quad \text{on } \mathcal{M}_T, \quad (2.13)$$

where N_T is a constant depending on K, d, d_1, T, λ .

We briefly outline how this is proved in Section 6. Following [1] we “shake” the finite difference scheme, we smooth the corresponding solution to get a supersolution of the normalized Bellman PDE (2.7). Hence we obtain the estimate $v \leq v_{\tau, h} + N(\tau^{1/4} + h^{1/2})$ by using a comparison of v with supersolutions to the normalized Bellman PDE. To get $v_{\tau, h} \leq v + N(\tau^{1/4} + h^{1/2})$ we use the same approach with the roles of v and $v_{\tau, h}$ interchanged. We “shake” the optimal control problem (2.5)-(2.6), again following [1]. We smooth the resulting payoff function to obtain a supersolution of the finite difference scheme. We get the estimate by using a comparison theorem established for the finite difference scheme.

An important consequence of Theorem 2.6 is that one gets the same rate of convergence for the optimal stopping and control problem.

Corollary 2.7. *Let w be the payoff function for the optimal stopping and control problem (2.3). If Assumptions 2.1, 2.3, 2.4 and 2.5 are satisfied with some positive function m^α , then*

$$\begin{aligned} & \sup_{\alpha \in A, r \geq 0} \left(m^\alpha \wedge \frac{1}{1+r} \right) [\delta_\tau u + L_h^\alpha u - ru + f^\alpha + rg] = 0 \quad \text{on } \mathcal{M}_T, \quad (2.14) \\ & u = g \quad \text{on } \bar{\mathcal{M}}_T \setminus \mathcal{M}_T, \end{aligned}$$

has a unique bounded solution $w_{\tau,h}$ and

$$|w - w_{\tau,h}| \leq N_T(\tau^{1/4} + h^{1/2}) \quad \text{on } \mathcal{M}_T,$$

where N_T is a constant depending on K, d, d_1, T, λ .

Proof. First we use Theorem 2.2 to see that $w(s, x)$ given by (2.3) is equal to (2.4). Consider a metric space $\bar{A} := A \times [0, \infty)$, with any metric defining the product topology. Let $\bar{A}_n = A_n \times [0, n]$. Then $\bar{A} = \bigcup_{n \in \mathbb{N}} \bar{A}_n$. Let $\bar{\mathfrak{A}}_n$ denote the set of progressively measurable processes taking values in $\bar{A}_n$. Clearly

$$w(s, x) = \sup_{\bar{\alpha} \in \bar{\mathfrak{A}}} \mathbb{E}_{s,x}^{\bar{\alpha}} \left[\int_0^{T-s} \bar{f}^{\bar{\alpha}_t}(s+t, x_t) e^{-\bar{\varphi}_t} dt + g(x_{T-s}) e^{-\bar{\varphi}_{T-s}} \right],$$

where

$$\bar{\mathfrak{A}} = \bigcup_{n \in \mathbb{N}} \bar{\mathfrak{A}}_n, \quad \bar{f}^{\bar{\alpha}} := f^\alpha + r g, \quad \bar{c}^{\bar{\alpha}} = r + c^\alpha, \quad \bar{\varphi}_t = \int_0^t \bar{c}^{\bar{\alpha}_u}(u, x_u) du$$

and x_t is the solution to (2.2). Notice that we are still using the same $\sigma^\alpha, \beta^\alpha$. We can check that Assumptions 2.1, 2.3, 2.4 and 2.5 hold with $\bar{A}$,

$$\bar{m}^{\bar{\alpha}} = \bar{m}^{(\alpha,r)} := m^\alpha \wedge \frac{1}{1+r},$$

$\bar{c}^\alpha$ and $\bar{f}^\alpha$ in place of A, m^α, f^α and c^α respectively. Finally we apply Theorem 2.6 to w and the scheme (2.14). $\square$

Corollary 2.8. *The system (2.14) is equivalent to:*

$$\begin{aligned} \sup_{\alpha \in A} m^\alpha [\delta_\tau u + L_h^\alpha u + f^\alpha] &< 0, \quad g - u = 0 \quad \text{on } \mathcal{M}_T, \\ \sup_{\alpha \in A} m^\alpha [\delta_\tau u + L_h^\alpha u + f^\alpha] &= 0, \quad g - u \leq 0 \quad \text{on } \mathcal{M}_T \\ \text{and } u &= g \quad \text{on } \bar{\mathcal{M}}_T \setminus \mathcal{M}_T. \end{aligned} \quad (2.15)$$

Proof. Let $\varepsilon = \frac{1}{1+r}$ in (2.14) and take the supremum over $\varepsilon \in [0, 1]$. Hence (2.14) can be rewritten as

$$\begin{aligned} \sup_{\varepsilon \in [0,1]} \left[\varepsilon \sup_{\alpha \in A} m^\alpha (\delta_\tau u + L_h^\alpha u + f^\alpha) + (1-\varepsilon)(g-u) \right] &= 0 \quad \text{on } \mathcal{M}_T, \\ u &= g \quad \text{on } \bar{\mathcal{M}}_T \setminus \mathcal{M}_T. \end{aligned}$$

Notice that the supremum over ε is achieved by either taking $\varepsilon = 0$ or $\varepsilon = 1$. Hence it can be seen that this is equivalent to (2.15). $\square$

3. ON THE SOLUTIONS OF THE FINITE DIFFERENCE SCHEME

First we give two simple examples which justify the condition (2.12).

Examples 3.1. Consider $A = [0, \infty)$, $m^\alpha = (1 + \alpha)^{-1}$ and the equation

$$\sup_{\alpha \in A} m^\alpha (\delta_\tau u) = 0 \quad \text{on } \mathcal{M}_T$$

with the terminal condition $u = 1$ on $\bar{\mathcal{M}}_T \setminus \mathcal{M}_T$. If $u : \mathcal{M}_T \rightarrow \mathbb{R}$ is any non-increasing function in t , then $m^\alpha \delta_\tau u \leq 0$. Hence, letting $\alpha \rightarrow \infty$, we

see that u satisfies the equation. Consequently the solution to the above problem is not unique.

Let $f^\alpha = 1 + \alpha$. Consider now the equation

$$\sup_{\alpha \in A} m^\alpha (\delta_\tau u + f^\alpha) = \sup_{\alpha \in A} m^\alpha \delta_\tau u + 1 = 0 \quad \text{on } \mathcal{M}_T.$$

If u is a solution then we have $m^\alpha \delta_\tau u \leq 0$. Hence $\sup_{\alpha \in A} m^\alpha \delta_\tau u = 0$, which contradicts the equation. Thus the above equation has no solution.

Let Assumption 2.4 be satisfied throughout the remainder of this section.

Lemma 3.2. *There is a unique bounded solution of the finite difference problem (2.9)-(2.10) on $\mathcal{M}_T$.*

Proof. Let $\gamma = (0, 1)$ and define ξ recursively as follows: $\xi(T) = 1$, $\xi(t) = \gamma^{-1}\xi(t + \tau_T(t))$ for $t < T$. Then for any function v

$$\delta_\tau(\xi v) = \gamma \xi \delta_\tau v - \nu \xi v, \quad \text{where } \nu = \frac{1 - \gamma}{\tau}.$$

To solve (2.9)-(2.10) for u , we could equivalently solve the following for v , with $u = \xi v$:

$$v = \mathbf{1}_{\mathcal{M}_T \setminus Q} \frac{1}{\xi} g + \mathbf{1}_Q G[v], \quad (3.1)$$

where for any $\varepsilon > 0$,

$$G[v] := v + \varepsilon \xi^{-1} \sup_{\alpha} m^\alpha (\delta_\tau u + L_h^\alpha u + f^\alpha).$$

Then, using the convention that repeated indices indicate summation and always summing up before taking the supremum,

$$G[v] = \sup_{\alpha} [p_\tau^\alpha T_\tau v + p_k^\alpha T_{h,l_k} v + p^\alpha v + \varepsilon m^\alpha (T_\tau \xi^{-1}) f^\alpha],$$

with

$$\begin{aligned} p_\tau^\alpha &= \varepsilon \gamma \tau^{-1} m^\alpha \geq 0, & p_k^\alpha &= \varepsilon (2h^{-2} a_k^\alpha + h^{-1} b_k^\alpha) m^\alpha \geq 0, \\ p^\alpha &= 1 - p_\tau^\alpha - \sum_k p_k^\alpha - \varepsilon \nu m^\alpha - \varepsilon m^\alpha c^\alpha. \end{aligned}$$

Notice that

$$p_k^\alpha \leq K \varepsilon \left(\frac{2K}{h^2} + \frac{K}{h} \right)$$

and

$$\varepsilon \nu m^\alpha + \varepsilon m^\alpha c^\alpha \leq \varepsilon \tau^{-1} K + \varepsilon K, \quad p_\tau \leq \varepsilon \tau^{-1} K,$$

so for all ε smaller than some ε_0 , $p^\alpha \geq 0$. Also

$$\begin{aligned} 0 \leq \sum_k p_k^\alpha + p^\alpha + p_\tau &= 1 - \varepsilon m^\alpha (\nu + c^\alpha) \leq 1 - (1 \vee \nu) \varepsilon m^\alpha (1 + c^\alpha) \\ &\leq 1 - (1 \vee \nu) \varepsilon K^{-1} =: \delta < 1, \end{aligned}$$

for some sufficiently small $\varepsilon > 0$. Since the difference of supremums is less than the supremum of a difference

$$|G[v](t, x) - G[w](t, x)| \leq \delta \sup_{\mathcal{M}_T} |v - w|.$$

Thus the operator G is a contraction on the space of bounded functions on $\bar{\mathcal{M}}_T$. By Banach's fixed point theorem (3.1) has a unique bounded solution. $\square$

Lemma 3.3. Consider f_1^α, f_2^α defined on $A \times \mathcal{M}_T$ and assume that in Q

$$\sup_{\alpha} m^\alpha f_2^\alpha < \infty, \quad f_1^\alpha \leq f_2^\alpha.$$

Let there be functions u_1, u_2 defined on $\bar{\mathcal{M}}_T$ and a constant $\mu \geq 0$ such that $u_1 e^{-\mu|x|}, u_2 e^{-\mu|x|}$ are bounded and for some $C \geq 0$

$$\begin{aligned} \sup_{\alpha} m^\alpha (\delta_\tau u_1 + L_h^\alpha u_1 + f_1^\alpha + C) \\ \geq \sup_{\alpha} m^\alpha (\delta_\tau u_2 + L_h^\alpha u_2 + f_2^\alpha) \quad \text{on } Q \end{aligned} \quad (3.2)$$

and

$$u_1 \leq u_2 \quad \text{on } \bar{\mathcal{M}}_T \setminus Q.$$

Then there exists a constant τ^* depending only on K, d_1, μ such for $\tau \in (0, \tau^*)$

$$u_1 \leq u_2 + TC \quad \text{on } \bar{\mathcal{M}}_T. \quad (3.3)$$

If u_1, u_2 are bounded on $\mathcal{M}_T$ then (3.3) holds for any τ .

To prove this lemma we use the following observation.

Remark 3.4. Let $D_\ell u := u_{x^i} \ell^i$, $D_\ell^2 u := u_{x^i x^j} \ell^i \ell^j$, $a_k^\alpha = \frac{1}{2}(\sigma_k^\alpha)^2$. Let D_x^n denote the collection of all n -th order derivatives in x . Consider

$$L^\alpha(t, x)u(t, x) = \sum_k a_k^\alpha(t, x) D_{\ell_k}^2 u(t, x) + b_k^\alpha(t, x) D_{\ell_k} u(t, x).$$

For any sufficiently smooth function $\eta(x)$, by Taylor's theorem,

$$m^\alpha |L^\alpha \eta(x) - L_h^\alpha \eta(x)| \leq N(h^2 \sup_{B_K(x)} |D_x^4 \eta| + h \sup_{B_K(x)} |D_x^2 \eta|).$$

Proof of Lemma 3.3. Let $w = u_1 - u_2 - C(T - t)$. From (3.2)

$$\sup_{\alpha} m^\alpha (\delta_\tau w + L_h^\alpha w) \geq 0$$

hence for any $\varepsilon > 0$

$$w + \varepsilon \sup_{\alpha} m^\alpha (\delta_\tau w + L_h^\alpha w) \geq w.$$

In the proof of Lemma 3.2, choose ε such that p^α, p_k^α are nonnegative. So the operator G is a monotone operator. Hence for any $\psi \geq w$,

$$\psi + \varepsilon \sup_{\alpha} m^\alpha (\delta_\tau \psi + L_h^\alpha \psi) \geq w. \quad (3.4)$$

Let $\gamma \in (0, 1)$. Use ξ from the proof of Lemma 3.2. Fix $\gamma \in (0, 1)$ later. Then

$$\delta_\tau \xi = \xi \frac{1}{\tau} (\gamma - 1).$$

Let

$$N_0 = \sup_{\bar{\mathcal{M}}_T} \frac{w_+}{\zeta} < \infty \quad \text{and} \quad \psi := N_0 \zeta \geq \zeta \frac{w_+}{\zeta} \geq w,$$

where $\eta(x) = \cosh(\mu|x|)$ and $\zeta = \eta\xi$. Then by Remark 3.4 and since for $r \geq 0$

$$\frac{\cosh(\mu r)}{r^2} - \frac{\sinh(\mu r)}{r^3} \leq \cosh(\mu r)$$

we get

$$m^\alpha L_h^\alpha \eta(x) \leq m^\alpha L^\alpha \eta(x) + N_1(h^2 + h) \cosh(\mu|x| + \mu K) \leq N_2 \cosh(\mu|x| + \mu K),$$

with N_1, N_2 depending only on μ, d_1, K . Hence

$$m^\alpha(\delta_\tau \zeta + L_h^\alpha \zeta) \leq \zeta[\tau^{-1}(\gamma - 1) + N_3], \quad N_3 := N_2 \frac{\cosh(\mu|x| + \mu K)}{\cosh(\mu|x|)} < \infty.$$

Then

$$\begin{aligned} w &\leq \psi + \varepsilon \sup_\alpha m^\alpha(\delta_\tau \psi + L_h^\alpha \psi) \\ &\leq N_0 \zeta [1 + \varepsilon(\tau^{-1}(\gamma - 1) + N_3)]. \end{aligned}$$

Define $\kappa(\gamma) = \tau^{-1}(\gamma - 1) + N_3$. Then for $\tau < \tau^* := N_3^{-1}$ one has $\kappa(0) < 0$ and $\kappa(1) > 0$. So one can have γ such that $\kappa < 0$ and $1 + \varepsilon\kappa > 0$. By (3.4)

$$w \leq \psi + \varepsilon \sup_\alpha m^\alpha(\delta_\tau \psi + L_h^\alpha \psi) \leq N_0 \xi(1 + \varepsilon\kappa) \quad \text{on } Q. \quad (3.5)$$

If $w \geq 0$ then $w_+ = w$. Hence

$$N_0 = \sup_{\mathcal{M}_T} \frac{w_+}{\xi} \leq \sup_{\mathcal{M}_T} \frac{N_0 \xi(1 + \varepsilon\kappa)}{\xi}. \quad (3.6)$$

By the hypothesis $w \leq 0$ on $\bar{\mathcal{M}}_T \setminus Q$. But $\kappa < 0$, hence N_0 must be 0 and so $w \leq 0$ on $\bar{\mathcal{M}}_T$. Finally observe that if $\mu = 0$, $N_2 = N_3 = 0$. $\square$

Corollary 3.5. *If v is the solution of (2.9)-(2.10) then*

$$v \leq 2K^2(T + 1) + \sup_x |g(x)|.$$

Proof. Apply Lemma 3.3 to v and

$$\xi(t) = K^2[(T - t) + 1] + N_1,$$

where $N_1 := \sup_x |g(x)|$. Then on Q

$$\begin{aligned} &\sup_\alpha m^\alpha(\delta_\tau \xi + L_h^\alpha \xi + f^\alpha) \\ &= \sup_\alpha [-m^\alpha(K^2 + c^\alpha(K^2[(T - t) + 1] + N_1))] \leq 0, \end{aligned}$$

while on $\bar{\mathcal{M}}_T \setminus Q$

$$\xi(T) \geq \sup_x |g(x)| \geq g(x).$$

$\square$

Corollary 3.6. *Assume that u_1, u_2 are functions both satisfying (2.9) on Q and $u_1 = g_1$, $u_2 = g_2$ on $\bar{\mathcal{M}}_T \setminus Q$. Then*

$$u_1 \leq u_2 + \sup_x (g_1 - g_2)_+.$$

Proof. Apply the comparison principle to u_1 and $\bar{u}_2 := \sup_x (g_1 - g_2)_+$. $\square$

Corollary 3.7. *Fix $(s_0, x_0) \in \bar{\mathcal{M}}_T$. Let v be a solution of (2.9)-(2.10). Define*

$$\nu = \sup_{(s_0, x) \in \bar{\mathcal{M}}_T} \frac{|v(s_0, x) - v(s_0, x_0)|}{|x - x_0|}.$$

Then for all $(t_0, x_0) \in \bar{\mathcal{M}}_T$ such that $t_0 \in [s_0 - 1, s_0]$,

$$|v(s_0, x_0) - v(t_0, x_0)| \leq N(\nu + 1)|s_0 - t_0|^{\frac{1}{2}}.$$

Proof. Assume, without loss of generality, that $\nu > 1$. Indeed if $\nu < 1$, then in the proof of the theorem use $\bar{\nu} = \nu + 1$. If the result holds for $\bar{\nu}$, then

$$N(\bar{\nu} + 1) \leq N(\nu + 1 + 1) \leq 2N(\nu + 1).$$

Assume that $s_0 > 0$. Shifting the time axis, so that $t_0 = 0$. Then $s_0 \leq 1$. Let

$$\gamma = s_0^{-1/2} \quad \text{and} \quad \xi(t) = \begin{cases} e^{s_0-t} & \text{for } t < s_0, \\ 1 & \text{for } t \geq s_0. \end{cases}$$

Notice that $\gamma \geq 1/T$. Define

$$\psi = \gamma\nu [\zeta + \kappa(s_0 - t)] + \frac{1}{\gamma}\nu\kappa + K(s_0 - t) + v(s_0, x_0),$$

where $\zeta = \eta\xi$, $\eta(x) = |x - x_0|^2$ and κ is a (large) constant to be chosen later, depending on K, d_1 and T only. The aim now is to apply Lemma 3.3 to v and ψ on $\mathcal{M}_{s_0}$.

$$\delta_\tau \xi(t) = -\theta\xi, \quad \text{where} \quad \theta = \tau^{-1}(1 - e^{-\tau}) \geq K^{-1}(1 - e^{-K}).$$

With $(\cdot, \cdot)$ denoting the inner product in $\mathbb{R}^d$

$$\mathbf{L}_h^\alpha \eta = 2a_k^\alpha |l_k|^2 + b_k^\alpha(l_k, 2(x - x_0) + hl_k) - c^\alpha \eta \leq N_1(1 + |x - x_0|) - c^\alpha \eta.$$

Hence

$$\begin{aligned} \delta_\tau \zeta + \mathbf{L}_h^\alpha \zeta &\leq \xi(t) [N_1(1 + |x - x_0|) - \theta|x - x_0|^2] - c^\alpha \zeta \\ &\leq N_2(1 + |x - x_0|) - \theta|x - x_0|^2 - c^\alpha \zeta. \end{aligned}$$

Since $v(s_0, x) \geq -K - \sup_x |g| \geq -N_3$ and hence

$$m^\alpha c^\alpha \psi \geq -m^\alpha c^\alpha \frac{\kappa}{\gamma} \nu + m^\alpha c^\alpha N_3.$$

Therefore

$$\begin{aligned} &m^\alpha (\delta_\tau \psi + \mathbf{L}_h^\alpha \psi + f^\alpha) \\ &\leq m^\alpha \left(\gamma\nu [N_2(1 + |x - x_0|) - \theta|x - x_0|^2 - \kappa] + c^\alpha N_3 - c^\alpha \frac{\kappa}{\gamma} \nu \right). \end{aligned} \quad (3.7)$$

The expression on the right hand side is a quadratic in $|x - x_0|$ with negative leading coefficient. It achieves its maximum of

$$m^\alpha \left\{ \gamma\nu \left[N_2 + \frac{1}{2\theta} - \frac{1}{4\theta N_2^2} - \kappa \right] - K \right\} + K + m^\alpha \left(c^\alpha N_3 - c^\alpha \frac{\kappa}{\gamma} \nu \right)$$

when $|x - x_0| = \frac{1}{2\theta N_2}$. Also $s_0 \leq 1$ and so $\gamma \geq 1$. Since $m^\alpha(1 + c^\alpha) \geq K^{-1}$, $m^\alpha(\gamma^2 + c^\alpha) \geq (1 \wedge \gamma^2)K^{-1}$. Hence

$$\kappa\nu\gamma^{-1}m^\alpha(\gamma^2 + c^\alpha) \geq \kappa\nu\gamma^{-1}(\gamma^2 \wedge 1)K^{-1} \geq K^{-1}\nu\kappa.$$

Finally, $m^\alpha c^\alpha N_T \leq K$. If κ is chosen very large (depending on K, T, d only) then (3.7) is non-positive. Hence

$$m^\alpha (\delta \psi + \mathbf{L}_h^\alpha \psi + f^\alpha) \leq 0.$$

To apply Lemma 3.3, $\psi \geq v$ on $\bar{\mathcal{M}}_{s_0} \setminus \mathcal{M}_{s_0}$ must be satisfied. Consider the two cases when either $\gamma|x - x_0| \leq 1$ or $\gamma|x - x_0| > 1$. Then

$$\begin{aligned} \psi(s_0, x) &= \nu [\gamma|x - x_0|^2 + \kappa\gamma^{-1}] + v(s_0, x_0) \\ &\geq \nu|x - x_0| + v(s_0, x_0) \geq v(s_0, x). \end{aligned}$$

By Lemma 3.3

$$v(t_0, x_0) - v(s_0, x_0) \leq \nu s_0^{-1/2} \kappa s_0 + s_0^{1/2} \nu \kappa + K s_0 \leq N(\nu + 1) s_0^{1/2}.$$

To get the estimate from the other side, one would consider

$$\psi = -\gamma \nu [\zeta + \kappa(s_0 - t)] - \frac{1}{\gamma} \nu \kappa - K(s_0 - t) + v(s_0, x_0)$$

and then show that for some choice of κ

$$\sup_{\alpha} m^{\alpha} (\delta \psi + L_h^{\alpha} \psi + f^{\alpha}) \geq 0 \quad \text{on } Q.$$

and that $\psi \leq v$ on $\bar{\mathcal{M}}_{s_0}$. $\square$

4. GRADIENT ESTIMATE FOR THE FINITE DIFFERENCE PROBLEM

The following lemma states some simple properties of the operators $\delta_{h,l}$ and $\Delta_{h,l}$, which will be used in this section. The statements of the following lemma are all proved in section 4 of [1].

Lemma 4.1. *Let u and v be functions on $\mathbb{R}^d$, l and x_0 vectors in $\mathbb{R}^d$ and $h > 0$. Then*

$$\delta_{h,l}(uv) = (\delta_{h,l} u) v + (T_{h,l} u) \delta_{h,l} v = v \delta_{h,l} u + u \delta_{h,l} v + h(\delta_{h,l} u)(\delta_{h,l} v), \quad (4.1)$$

which can be thought of as the discrete Leibnitz rule. Also

$$\begin{aligned} \Delta_{h,l}(uv) &= v \Delta_{h,l} u + (\delta_{h,l} u) \delta_{h,l} v + (\delta_{h,-l} u) \delta_{h,-l} v + u \Delta_{h,l} v, \\ \Delta_{h,l}(u^2) &= 2u \Delta_{h,l} u + (\delta_{h,l} u)^2 + (\delta_{h,-l} u)^2. \end{aligned} \quad (4.2)$$

Furthermore if $v(x_0) \leq 0$ then at x_0 it holds that

$$-\delta_{h,l} v \leq \delta_{h,l}(v_-), \quad -\Delta_{h,l} v \leq \Delta_{h,l}(v_-), \quad (4.3)$$

$$-\delta_{h,l}(u_-) \leq [\delta_{h,\ell}((u+v)_-)]_- + [\delta_{h,\ell}(v_-)]_+. \quad (4.4)$$

$$\begin{aligned} (\Delta_{h,\ell} u)_- &\leq [\delta_{h,-\ell}((\delta_{h,\ell} u + v)_-)]_- + [\delta_{h,\ell}((\delta_{h,-\ell} u + w)_-)]_- \\ &\quad + [\delta_{h,-\ell}(v_-)]_+ + [\delta_{h,\ell}(w_-)]_+. \end{aligned} \quad (4.5)$$

$$|\Delta_{h,l} u| \leq |\delta_{h,-l}((\delta_{h,l} u)_-)| + |\delta_{h,l}((\delta_{h,-l} u)_-)|, \quad (4.6)$$

$$|\Delta_{h,l} u| \leq |\delta_{h,-l}((\delta_{h,l} u)_+)| + |\delta_{h,l}((\delta_{h,-l} u)_+)|, \quad (4.7)$$

Let Assumption 2.4 hold. Let T' be the smallest integer multiple of τ which is greater than or equal to T . Choose an arbitrary $\varepsilon \in (0, Kh]$ and $l \in R^d$. Let $h_r = h$ for $r = \pm 1, \dots, \pm d_1$ and $h_r = \varepsilon$ for $r = \pm(d_1 + 1)$. Let

$$\bar{\mathcal{M}}_T(\varepsilon) := \{(t, x + i\varepsilon l) : (t, x) \in \bar{\mathcal{M}}_T, i = 0, \pm 1, \dots\}.$$

For a general $Q \subset \bar{\mathcal{M}}_T(\varepsilon)$ let

$$\begin{aligned} Q_{\varepsilon}^0 &= \{(t, x) : (t + \tau_T(t), x) \in Q, (t, x + h_r l_r) \in Q, \forall r = \pm 1, \dots, \pm(d_1 + 1)\}, \\ \partial_{\varepsilon} Q &= Q \setminus Q_{\varepsilon}^0. \end{aligned}$$

Assumption 4.2. Assume that for any $(t, x) \in Q_{\varepsilon}^0$

$$|\delta_{h_r, l_r} b_k^{\alpha}| \leq K, \quad m^{\alpha} |\delta_{h_r, l_r} f^{\alpha}| \leq K, \quad |\delta_{h_r, l_r} c^{\alpha}| \leq K, \quad (4.8)$$

$$|\delta_{h_r, l_r} a_k^{\alpha}| \leq K \sqrt{a_k^{\alpha}} + h. \quad (4.9)$$

Define $a^\pm = a_\pm = (1/2)(|a| \pm a)$.

Theorem 4.3. *Let u be a function defined on $\bar{\mathcal{M}}_T(\varepsilon)$ satisfying (2.9) on Q . If Assumption 4.2 is satisfied, then there is a constant $N > 1$, such that, if there exists some $c_0 \geq 0$ satisfying*

$$\lambda + \frac{1}{\tau}(1 - e^{-c_0\tau}) > N, \quad (4.10)$$

then on Q ,

$$|\delta_{\varepsilon, \pm l} u| \leq N e^{c_0 T'} (1 + |u|_{0,Q} + \sup_{k, \partial_\varepsilon Q} (|\delta_{h_k, \ell_k} u| + |\delta_{\varepsilon, l} u| + |\delta_{\varepsilon, -l} u|)). \quad (4.11)$$

Proof of Theorem 4.3. Let

$$v_r = \delta_{h_r, l_r} v, \quad v = \xi u, \quad \xi(t) = \begin{cases} e^{c_0 t} & t < T, \\ e^{c_0 T'} & t = T. \end{cases}$$

Let $(t_0, x_0) \in Q$ be the point where

$$V = \sum_r (v_r^-)^2$$

is maximized. By definition, for any $(t, x) \in Q_\varepsilon^0$ we know that

$$(t, x + h_r l_r) \in Q.$$

Then either

$$v_r(t, x) \leq 0 \quad \text{or} \quad -v_r(t, x) = v_{-r}(t, x + h_r l_r) \leq 0.$$

In either case, for any $(t, x) \in Q_\varepsilon^0$

$$|v_r(t, x)| \leq V^{1/2}.$$

Hence

$$M_1 := \sup_{Q, r} |v_r| \leq \sup_{\partial_\varepsilon Q, r} |v_r| + V^{1/2},$$

$$|\delta_{\varepsilon, \pm l} u| \leq e^{c_0 T'} \sup_{\partial_\varepsilon Q, r} |\delta_{h_r, l_r} u| + V^{1/2}$$

on Q . So we only have to estimate V on Q . If (t_0, x_0) belongs to $\partial_\varepsilon Q$, then the conclusion of the theorem is trivially true. Thus, we may assume that $(t_0, x_0) \in Q_\varepsilon^0$. Then for any $\bar{\varepsilon} > 0$ there exists $\alpha_0 \in A$ such that at (t_0, x_0) ,

$$m^{\alpha_0} (\delta_\tau u + a_k^{\alpha_0} \Delta_{h_k, \ell_k} u + b_k^{\alpha_0} \delta_{h_k, \ell_k} u - c^{\alpha_0} u + f^{\alpha_0}) + \bar{\varepsilon} \geq 0.$$

and so for some $\bar{\varepsilon} \in (0, \bar{\varepsilon}_0]$

$$m^{\alpha_0} (\delta_\tau u + a_k^{\alpha_0} \Delta_{h_k, \ell_k} u + b_k^{\alpha_0} \delta_{h_k, \ell_k} u - c^{\alpha_0} u + f^{\alpha_0}) + \bar{\varepsilon} = 0. \quad (4.12)$$

Furthermore (thanks to the fact that $(t_0, x_0) \in Q_\varepsilon^0$)

$$T_{h_r, \ell_r} [m^{\alpha_0} (\delta_\tau u + a_k^{\alpha_0} \Delta_{h_k, \ell_k} u + b_k^{\alpha_0} \delta_{h_k, \ell_k} u - c^{\alpha_0} u + f^{\alpha_0})] \leq 0. \quad (4.13)$$

We subtract (4.12) from (4.13) and divide by h_r to obtain that for each r

$$m^{\alpha_0} \delta_{h_r, l_r} (\delta_\tau u + a_k^{\alpha_0} \Delta_{h_k, \ell_k} u + b_k^{\alpha_0} \delta_{h_k, \ell_k} u + f^{\alpha_0} - c^{\alpha_0} u) - \frac{\bar{\varepsilon}}{h_r} \leq 0.$$

For each r , by the discrete Leibnitz rule (4.1)

$$\begin{aligned} m^{\alpha_0} (\delta_\tau (\xi^{-1} v_r) + \xi^{-1} [a_k^{\alpha_0} \Delta_{h_k, \ell_k} v_r + I_{1r} + I_{2r} + I_{3r}] + \delta_{h_r, l_r} f^{\alpha_0}) \\ - \xi^{-1} \delta_{h_r, l_r} (m^{\alpha_0} c^{\alpha_0} v) - \frac{\bar{\varepsilon}}{h_r} \leq 0, \end{aligned} \quad (4.14)$$

where

$$\begin{aligned} I_{1r} &= (\delta_{h_r, l_r} a_k^{\alpha_0}) \Delta_{h_k, \ell_k} v, \\ I_{2r} &= h_r (\delta_{h_r, l_r} a_k^{\alpha_0}) \Delta_{h_k, \ell_k} v_r, \\ I_{3r} &= (T_{h_r, \ell_r} b_k^{\alpha_0}) \delta_{h_k, \ell_k} v_r + (\delta_{h_r, l_r} b_k^{\alpha_0}) \delta_{h_k, \ell_k} v. \end{aligned}$$

Notice (first equality by (4.2), second inequality by (4.3), also note that there is a tacit summation in k) that

$$\begin{aligned} 0 &\geq \Delta_{h_k, \ell_k} \sum_r (v_r^-)^2 = 2v_r^- \Delta_{h_k, \ell_k} v_r^- + \sum_r [\delta_{h_k, \ell_k} (v_r^-)^2 + \delta_{h_k, l_{-k}} (v_r^-)^2] \\ &\geq -2v_r^- \Delta_{h_k, \ell_k} v_r + 2 \sum_r (\delta_{h_k, \ell_k} v_r^-)^2. \end{aligned}$$

Since a_k^α are nonnegative,

$$0 \leq v_r^- \Delta_{h_k, \ell_k} v_r \quad (4.15)$$

and

$$I := \sum_r a_k^{\alpha_0} (\delta_{h_k, \ell_k} v_r^-)^2 \leq v_r^- a_k^{\alpha_0} \Delta_{h_k, \ell_k} v_r.$$

Go back to (4.14), multiply by ξv_r^- and sum up in r to get

$$\begin{aligned} m^{\alpha_0} \left(\xi v_r^- \delta_\tau (\xi^{-1} v_r) + \frac{1}{2} a_k^{\alpha_0} v_r^- \Delta_{h_k, \ell_k} v_r + \frac{1}{2} I + v_r^- [I_{1r} + I_{2r} \right. \\ \left. + I_{3r} + \xi \delta_{h_r, l_r} f^{\alpha_0}] \right) - v_r^- \delta_{h_r, l_r} (m^{\alpha_0} c^{\alpha_0} v) - \sum_r \xi (v_r^-) \bar{\bar{\xi}} \leq 0. \end{aligned} \quad (4.16)$$

By (4.8), $m^\alpha \delta_{h_r, l_r} f_k^\alpha \geq -K$. If $v_r > 0$, then $v_r^- = 0$, hence $-v_r^- v_r = 0 = (v_r^-)^2$. On the other hand if $v_r \leq 0$, then $v_r = -v_r^-$ and so $-v_r^- v_r = (v_r^-)^2$. Hence

$$\begin{aligned} m^{\alpha_0} v_r^- \xi \delta_{h_r, l_r} f^{\alpha_0} - v_r^- \delta_{h_r, l_r} (m^{\alpha_0} c^{\alpha_0} v) \\ = m^{\alpha_0} v_r^- \xi \delta_{h_r, l_r} f^{\alpha_0} - m^{\alpha_0} v_r^- (\delta_{h_r, l_r} c^{\alpha_0}) v - m^{\alpha_0} v_r^- (T_{h_r, \ell_r} c^{\alpha_0}) v_r \\ \geq -e^{c_0 T'} K v_r^- - m^{\alpha_0} v_r^- |\delta_{h_r, l_r} c^{\alpha_0}| |v| + m^{\alpha_0} (T_{h_r, \ell_r} c^{\alpha_0}) (v_r^-)^2 \\ \geq -e^{c_0 T'} K M_1 - m^{\alpha_0} K M_1 M_o + (T_{h_r, \ell_r} m^{\alpha_0} c^{\alpha_0}) V. \end{aligned}$$

Use the fact that V attains its maximum at (t_o, x_o) , discrete Leibnitz rule and (4.3) in order to get a lower bound for the term containing I_{3r} . First,

$$\begin{aligned} 0 &\geq \sum_r \delta_{h_k, \ell_k} (v_r^-)^2 = 2v_r^- \delta_{h_k, \ell_k} v_r^- + \sum_r h_k (\delta_{h_k, \ell_k} v_r^-)^2 \\ &\geq 2v_r^- \delta_{h_k, \ell_k} v_r^- \geq -2v_r^- \delta_{h_k, \ell_k} v_r. \end{aligned}$$

Next recall that $b_k^\alpha \geq 0$ and $|\delta_{h_r, l_r} b_k^\alpha| \leq K$, and so

$$-v_r^- (T_{h_r, \ell_r} b_k^{\alpha_0}) \delta_{h_k, \ell_k} v_r \leq 0,$$

hence

$$v_r^- I_{3r} \geq -v_r^- |\delta_{h_r, l_r} b_k^{\alpha_0}| |\delta_{h_k, \ell_k} v| \geq -K M_1^2.$$

Apply discrete Leibnitz rule to the very first term of (4.16). Then

$$\begin{aligned} \xi v_r^- \delta_\tau (\xi^{-1} v_r) &= \xi v_r^- [\xi^{-1} (t_o + \tau) \delta_\tau v_r + v_r \delta_\tau \xi^{-1}] \\ &= e^{-c_0 \tau} v_r^- \delta_\tau v_r - V \xi \delta_\tau \xi^{-1} \geq -V \xi \delta_\tau \xi^{-1} \\ &= V \frac{1}{\tau} (1 - e^{-c_0 \tau}). \end{aligned}$$

Using the above estimates we see

$$\begin{aligned} m^{\alpha_0} V \frac{1}{\tau} (1 - e^{-c_0 \tau}) + (T_{h_r, \ell_r} m^{\alpha_0} c^{\alpha_0}) V + \frac{1}{2} m^{\alpha_0} a_k^{\alpha_0} v_r^- \Delta_{h_k, \ell_k} v_r + \frac{1}{2} m^{\alpha_0} I \\ + v_r^- m^{\alpha_0} [I_{1r} + I_{2r}] - \sum_r \xi v_r^- \frac{\bar{\varepsilon}}{h_r} \leq K M_1 (e^{c_0 T'} + M_0 + m^{\alpha_0} M_1). \end{aligned}$$

Hence

$$\begin{aligned} m^{\alpha_0} V \frac{1}{\tau} (1 - e^{-c_0 \tau}) + (T_{h_r, \ell_r} m^{\alpha_0} c^{\alpha_0}) V \leq K M_1 (e^{c_0 T'} + M_0 + m^{\alpha_0} M_1) \\ + m^{\alpha_0} v_r^- |I_{1r}| + m^{\alpha_0} v_r^- |I_{2r}| - \frac{1}{2} m^{\alpha_0} a_k^{\alpha_0} v_r^- \Delta_{h_k, \ell_k} v_r - \frac{1}{2} m^{\alpha_0} I + \sum_r \xi v_r^- \frac{\bar{\varepsilon}}{h_r}. \end{aligned}$$

Define

$$\begin{aligned} J_1 &:= v_r^- |(\delta_{h_r, \ell_r} a_k^{\alpha_0}) \Delta_{h_k, \ell_k} v| - \frac{1}{4} \sum_r a_k^{\alpha_0} (\delta_{h_k, \ell_k} v_r^-)^2, \\ J_2 &:= J_3 - \frac{1}{2} a_k^{\alpha_0} v_r^- \Delta_{h_k, \ell_k} v_r - \frac{1}{4} \sum_r a_k^{\alpha_0} (\delta_{h_k, \ell_k} v_r^-)^2 \\ J_3 &:= h_r v_r^- |(\delta_{h_r, \ell_r} a_k^{\alpha_0}) \Delta_{h_k, \ell_k} v_r|. \end{aligned}$$

One can rewrite the above inequality as

$$\begin{aligned} m^{\alpha_0} V \frac{1}{\tau} (1 - e^{-c_0 \tau}) + (T_{h_r, \ell_r} m^{\alpha_0} c^{\alpha_0}) V \\ \leq K M_1 (e^{c_0 T'} + M_0 + m^{\alpha_0} M_1) + m^{\alpha_0} (J_1 + J_2) + \sum_r \xi v_r^- \frac{\bar{\varepsilon}}{h_r}. \end{aligned}$$

So we need to estimate J_1, J_2 . By (4.6)

$$|\Delta_{h_k, \ell_k} v| \leq \sum_r |\delta_{h_k, \ell_k} v_r^-| + \sum_r |\delta_{h_k, \ell_{-k}} v_r^-|$$

then because there's tacit summation in $k = \pm 1, \dots, \pm d_1$ and because in the second sum we can reverse the order of summation in k

$$|\Delta_{h_k, \ell_k} v| \leq 2 \sum_r |\delta_{h_k, \ell_k} v_r^-|.$$

We turn our attention to J_1 .

$$\begin{aligned} v_r^- |(\delta_{h_r, \ell_r} a_k^{\alpha_0}) \Delta_{h_k, \ell_k} v| &\leq M_1 K (\sqrt{a_k^{\alpha_0}} + h) |\Delta_{h_k, \ell_k} v| \\ &\leq K M_1 \sqrt{a_k^{\alpha_0}} |\Delta_{h_k, \ell_k} v| + K M_1^2 \\ &\leq K M_1^2 + 2 K M_1 \sqrt{a_k^{\alpha_0}} \sum_r \delta_{h_k, \ell_k} v_r^- \\ &\leq K M_1^2 + \sum_r 8 K M_1^2 + \frac{1}{4} \sum_r a_k^{\alpha_0} (\delta_{h_k, \ell_k} v_r^-)^2, \end{aligned}$$

having used Young's inequality. So $J_1 \leq N M_1^2$. Since $h_r = h$ or $\varepsilon \in (0, Kh]$,

$$J_3 \leq K h v_r^- \sqrt{a_k^{\alpha_0}} |\Delta_{h_k, \ell_k} v_r| + K h^2 v_r^- |\Delta_{h_k, \ell_k} v_r|.$$

Also $h^2 |\Delta_{h_k, \ell_k} v_r| \leq 4 M_1$ and in general $|a| = 2a^- + a$ and so

$$J_3 \leq 2 K h v_r^- \sqrt{a_k^{\alpha_0}} (\Delta_{h_k, \ell_k} v_r)_- + K h v_r^- \sqrt{a_k^{\alpha_0}} |\Delta_{h_k, \ell_k} v_r| + N M_1^2.$$

By (4.5) (with summation in r everywhere)

$$h (\Delta_{h_k, \ell_k} v_r)_- \leq h |\Delta_{h_k, \ell_k} (v_r^-)| \leq |\delta_{h_k, \ell_k} (v_r^-)| + |\delta_{h_k, \ell_{-k}} (v_r^-)|.$$

Therefore

$$\begin{aligned}
J_3 &\leq 2K v_r^- h \sqrt{a_k^{\alpha_0}} \Delta_{h_k, \ell_k} v_r + 4KM_1 \left(\sum_r a_k^{\alpha_0} (\delta_{h_k, \ell_k} v_r^-)^2 \right)^{1/2} + NM_1^2 \\
&\leq NM_1^2 + 2K v_r^- h \sqrt{a_k^{\alpha_0}} \Delta_{h_k, \ell_k} v_r + \frac{1}{4} \sum_r a_k^{\alpha_0} (\delta_{h_k, \ell_k} v_r^-)^2, \\
J_2 &\leq NM_1^2 - \frac{1}{2} (a_k^{\alpha_0} - 4Kh \sqrt{a_k^{\alpha_0}}) v_r^- \Delta_{h_k, \ell_k} v_r.
\end{aligned}$$

Consider $\mathcal{K} := \{k : a_k^{\alpha_0} - 4Kh \sqrt{a_k^{\alpha_0}} \geq 0\}$. For all k not in $\mathcal{K}$ we have $a_k^{\alpha_0} < Nh^2$. Also $h^2 |\Delta_{h_k, \ell_k} v_r| \leq 4M_1$. Hence

$$\begin{aligned}
J_2 &\leq NM_1^2 \quad \text{and} \\
m^{\alpha_0} V \frac{1}{\tau} (1 - e^{-c_0 \tau}) + (T_{h_r, \ell_r} m^{\alpha_0} c^{\alpha_0}) V \\
&\leq NM_1 \left(e^{c_0 T'} + M_0 + m^{\alpha_0} M_1 \right) + \sum_r \xi v_r^- \frac{\bar{\varepsilon}}{h_r}.
\end{aligned}$$

We know from (5.5), that $M_1 \leq \mu + V^{1/2}$, where

$$\mu := \sup_{\partial_\varepsilon Q, r} |v_r| \leq e^{c_0 T'} \sup_{\partial_\varepsilon Q, r} |\delta_{h_r, \ell_r} u| =: \bar{\mu}.$$

One also defines

$$\bar{M}_0 = |u|_{0, Q} \geq e^{-c_0 T'} M_0, \quad \bar{V} = e^{-2c_0 T'} V$$

and so, using Young's inequality

$$\begin{aligned}
m^{\alpha_0} V \frac{1}{\tau} (1 - e^{-c_0 \tau}) + (T_{h_r, \ell_r} m^{\alpha_0} c^{\alpha_0}) V \\
&\leq N(\bar{\mu} + \bar{V}^{1/2}) \left(1 + \bar{M}_0 + \bar{\mu} + m^{\alpha_0} \bar{V}^{1/2} \right) + \sum_r \xi v_r^- \frac{\bar{\varepsilon}}{h_r} \\
&\leq N^* \left(1 + \bar{M}_0^2 + \bar{\mu}^2 + m^{\alpha_0} \bar{V} \right) + \sum_r \xi v_r^- \frac{\bar{\varepsilon}}{h_r}. \quad (4.17)
\end{aligned}$$

By assumption (5.3), we can find c_0 such that

$$\frac{1}{\tau} (1 - e^{-c_0 \tau}) > 1 + N^*.$$

Using this and $\bar{V} \leq V$, (4.17) becomes

$$m^{\alpha_0} V (1 + T_{h_r, \ell_r} c^{\alpha_0}) \leq N^* \left(1 + \bar{M}_0^2 + \bar{\mu}^2 \right) + \sum_r \xi v_r^- \frac{\bar{\varepsilon}}{h_r}.$$

By our assumptions on m^α and c^α , $m^{\alpha_0} (1 + T_{h_r, \ell_r} c^\alpha) \geq K^{-1}$ and so

$$V \leq KN^* \left(1 + \bar{M}_0^2 + \bar{\mu}^2 \right) + K \sum_r \xi v_r^- \frac{\bar{\varepsilon}}{h_r}.$$

Noticing that $\xi v_r^- \leq e^{c_0 T'} 2M_0 h_r^{-1}$

$$V \leq N^* \left(1 + \bar{M}_0^2 + \bar{\mu}^2 \right) + e^{c_0 T'} N M_0 \frac{\bar{\varepsilon}}{h_r^2}.$$

Neither N , nor M_0 depend on choice of α_0 and so we can let $\bar{\varepsilon} \rightarrow 0$. Then

$$V \leq N^* \left(1 + \bar{M}_0^2 + \bar{\mu}^2 \right).$$

□

Theorem 4.3 can now be used to estimate the difference between two solutions to the finite difference problem, based on how much the respective drift and diffusion coefficients, discounting factor and reward functions differ on the domain Q .

Remark 4.4. For small $\tau > 0$, (4.10) is satisfied. Indeed, for $\tau^{-1} > 2N$, where N is the constant arising in (4.11) let $c_0 \geq 0$ satisfy $e^{-c_0\tau} \leq 1/2$. Then c_0 depends on K, d_1 only.

Theorem 4.5. *Let*

$$\varepsilon = \sup_{\mathcal{M}_T, A, k} \left(|\sigma_k^\alpha - \hat{\sigma}_k^\alpha| + |b_k^\alpha - \hat{b}_k^\alpha| + |c^\alpha - \hat{c}^\alpha| + m^\alpha |f^\alpha - \hat{f}^\alpha| \right).$$

Assume that the functions $u, \hat{u}$ satisfy (2.9) on $Q = \mathcal{M}_T$, with the coefficients σ, b, c, f and $\hat{\sigma}, \hat{b}, \hat{c}, \hat{f}$ respectively. Then

$$|u - \hat{u}| \leq N e^{c_0 T'} T' \varepsilon I, \quad (4.18)$$

where

$$I := \sup_{(T, x) \in \mathcal{M}_T} \left(1 + \max_k (|\delta_{h_k, \ell_k} u| + |\delta_{h_k, \ell_k} \hat{u}| + \frac{1}{\varepsilon} |u(T, x) - \hat{u}(T, x)|) \right). \quad (4.19)$$

To prove the theorem we need the following lemma. Recall that $a_k^\alpha = (1/2)(\sigma_k^\alpha)^2$.

Lemma 4.6. *Let u and $\hat{u}$ be functions satisfying (2.9) on some $Q \subset \mathcal{M}_T$ with the coefficients σ, b, c, f and $\hat{\sigma}, \hat{b}, \hat{c}, \hat{f}$ respectively. Assume that $\varepsilon \in (0, h]$ and for all $\alpha \in A$*

$$\begin{aligned} |b_k^\alpha - \hat{b}_k^\alpha| + m^\alpha |f^\alpha - \hat{f}^\alpha| + |c^\alpha - \hat{c}^\alpha| &\leq \varepsilon, \\ |a_k^\alpha - \hat{a}_k^\alpha| &\leq 2\varepsilon \sqrt{a_k^\alpha \wedge \hat{a}_k^\alpha} + \varepsilon h \quad \text{on } Q. \end{aligned}$$

Then on Q

$$\begin{aligned} &|u - \hat{u}| \\ &\leq \varepsilon N e^{c_0 T'} \left[1 + |u|_{0, Q} + |\hat{u}|_{0, Q} + \sup_{k, \partial Q} (|\delta_{h_k, \ell_k} u| + |\delta_{h_k, \ell_k} \hat{u}| + \varepsilon^{-1} |u - \hat{u}|) \right]. \end{aligned}$$

We can deduce this from the gradient estimate by introducing an extra coordinate to the grid and by defining a suitable equation on the extended grid.

Proof. We will work with $(t, x) = (t, x', x^{d+1}) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}$. Let

$$\tilde{Q} := Q \times \{0, \pm\varepsilon, \dots, \pm m\varepsilon\},$$

$$\begin{aligned} \tilde{Q}_\varepsilon^0 &:= \{(t, x', x^{d+1}) : (t + \tau_T(t), x', x^{d+1}) \in \tilde{Q}, (t, x' + h\ell_k, x^{d+1} \pm \varepsilon) \in \tilde{Q}, \\ &\quad \forall k = 0, \pm 1, \dots, \pm d_1\} \end{aligned}$$

$$\partial_\varepsilon \tilde{Q} := \tilde{Q} \setminus \tilde{Q}_\varepsilon^0 = \partial Q \times \{0, \pm\varepsilon, \dots, \pm m\varepsilon\} \cup Q^0 \times \{m\varepsilon, -m\varepsilon\}.$$

where $m \geq \varepsilon^{-1}$ is a fixed integer. Also let

$$\tilde{a}_k^\alpha(t, x', x^{d+1}) = \begin{cases} a_k^\alpha(t, x') & \text{if } x^{d+1} > 0, \\ \hat{a}_k^\alpha(t, x') & \text{if } x^{d+1} \leq 0 \end{cases}$$

and define $\tilde{b}_k^\alpha, \tilde{c}_k^\alpha$ in a similar way. Let

$$\tilde{f}_k^\alpha(t, x', x^{d+1}) = \begin{cases} f_k^\alpha(t, x') \left[1 - \frac{x^{d+1} - \varepsilon}{\varepsilon m}\right] & \text{if } x^{d+1} > 0, \\ \hat{f}_k^\alpha(t, x') \left[1 + \frac{x^{d+1}}{\varepsilon m}\right] & \text{if } x^{d+1} \leq 0 \end{cases}$$

and define $\tilde{u}$ analogously. Then $\tilde{u}$ satisfies (2.9) with $\tilde{a}_k^\alpha, \tilde{b}_k^\alpha, \tilde{c}^\alpha, \tilde{f}^\alpha$ in place of $a_k^\alpha, b_k^\alpha, c^\alpha, f^\alpha$ respectively and with the normalizing factor m^α , on the domain $\tilde{Q}$. To apply Theorem 4.3 to $\tilde{u}$ one needs to ascertain that Assumption 4.2 is satisfied. For $r = \pm 1, \dots, \pm d_1$ (4.8) follows from Assumption 2.4. Furthermore, for $r = \pm(d_1 + 1)$ notice that

$$\delta_{\varepsilon, \ell} \tilde{a}_k^\alpha(t, x) = \begin{cases} 0 & \text{if } x^{d+1} > 0, \\ \varepsilon^{-1}(a_k^\alpha(t, x') - \hat{a}_k^\alpha(t, x')) & \text{if } x^{d+1} = 0, \\ 0 & \text{if } x^{d+1} < 0 \end{cases}$$

and similarly for $\tilde{b}_k^\alpha, \tilde{c}^\alpha$. So for $\tilde{b}_k^\alpha$ and $\tilde{c}^\alpha$ (4.8) holds, while (4.9) follows from:

$$\varepsilon^{-1} |a_k^\alpha(t, x') - \hat{a}_k^\alpha(t, x')| \leq K \sqrt{\tilde{a}_k^\alpha(t, x', 0)} + Kh.$$

Finally

$$m^\alpha \delta_{\varepsilon, \ell} \tilde{f}^\alpha(t, x) = \begin{cases} -m^\alpha f^\alpha(t, x') / (\varepsilon m) & \text{if } x^{d+1} > 0, \\ \varepsilon^{-1} m^\alpha (f^\alpha(t, x') - \hat{f}_k^\alpha(t, x')) & \text{if } x^{d+1} = 0, \\ m^\alpha \hat{f}^\alpha(t, x') / (\varepsilon m) & \text{if } x^{d+1} < 0 \end{cases}$$

shows (4.8) for $\tilde{f}^\alpha$. By Theorem 4.3, for all $(t, x') \in \mathcal{M}_T$

$$\begin{aligned} \varepsilon^{-1} |u(t, x') - \hat{u}(t, x')| &= |\delta_{\varepsilon, \ell} \tilde{u}(t, x', 0)| \leq NT e^{c_0(T+\tau)} \\ &\times \left[1 + |\tilde{u}|_{0, \tilde{Q}} + \sup_{\partial_\varepsilon \tilde{Q}} \left(\max_k |\delta_{h_k, \ell_k} \tilde{u}| + |\delta_{\varepsilon, \ell} \tilde{u}| + |\delta_{\varepsilon, -\ell} \tilde{u}| \right) \right], \end{aligned} \quad (4.20)$$

where N is, in particular, independent of m . To complete the proof consider $\partial_\varepsilon \tilde{Q}$. Either $(t, x', x^{d+1}) \in \partial Q \times 0, \pm\varepsilon, \dots, \pm m\varepsilon$. Then

$$\delta_{h_k, \ell_k} \tilde{u}(t, x', x^{d+1}) = \begin{cases} \delta_{h_k, \ell_k} u(t, x') \left[1 - \frac{x^{d+1} - \varepsilon}{\varepsilon m}\right] & \text{if } x^{d+1} > 0, \\ \delta_{h_k, \ell_k} \hat{u}(t, x') \left[1 + \frac{x^{d+1}}{\varepsilon m}\right] & \text{if } x^{d+1} \leq 0. \end{cases}$$

And so $|\delta_{h_k, \ell_k} \tilde{u}(t, x', x^{d+1})| \leq |\delta_{h_k, \ell_k} u(t, x')| + |\delta_{h_k, \ell_k} \hat{u}(t, x')|$. Also notice

$$\begin{aligned} \delta_{\varepsilon, \ell} \tilde{u}(t, x', x^{d+1}) &= \begin{cases} 0 & \text{if } x^{d+1} \neq 0, \\ \varepsilon^{-1} (u(t, x') - \hat{u}(t, x')) & \text{if } x^{d+1} = 0. \end{cases} \\ \delta_{\varepsilon, -\ell} \tilde{u}(t, x', x^{d+1}) &= \begin{cases} 0 & \text{if } x^{d+1} \neq 0, \\ \varepsilon^{-1} (\hat{u}(t, x') - u(t, x')) & \text{if } x^{d+1} = 0. \end{cases} \end{aligned}$$

Or $(t, x', x^{d+1}) \in Q^0 \times \{m\varepsilon, -m\varepsilon\}$. Then

$$\delta_{h_k, \ell_k} \tilde{u}(t, x', x^{d+1}) = \begin{cases} \delta_{h_k, \ell_k} u(t, x') \left[1 - \frac{\varepsilon m - \varepsilon}{\varepsilon m}\right] & \text{if } x^{d+1} = m\varepsilon, \\ \delta_{h_k, \ell_k} \hat{u}(t, x') \left[1 + \frac{-\varepsilon m}{\varepsilon m}\right] & \text{if } x^{d+1} = -m\varepsilon. \end{cases}$$

Notice that $[1 - \frac{\varepsilon m - \varepsilon}{\varepsilon m}] = 1/m$ and $[1 + \frac{-\varepsilon m}{\varepsilon m}] = 0$. Since $m \geq \varepsilon^{-1}$, $m\varepsilon \geq 1$ and so for $(t, x', x^{d+1}) \in Q^0 \times \{m\varepsilon, -m\varepsilon\}$

$$\delta_{\varepsilon, \ell} \tilde{u} = \delta_{\varepsilon, -\ell} \tilde{u} = 0.$$

Hence

$$\begin{aligned} & \sup_{\partial_\varepsilon \tilde{Q}} \left(\max_k |\delta_{h_k, \ell_k} \tilde{u}| + |\delta_{\varepsilon, \ell} \tilde{u}| + |\delta_{\varepsilon, -\ell} \tilde{u}| \right) \\ & \leq \sup_{k, \partial Q} \left(|\delta_{h_k, \ell_k} u| + 2|\delta_{h_k, \ell_k} \hat{u}| + 2\varepsilon^{-1}|u - \hat{u}| + |\delta_{h_k, \ell_k} u| \frac{1}{m} \right). \end{aligned}$$

Since the constant N in (4.20) is independent of m , we can let $m \rightarrow \infty$ and the conclusion of the lemma follows. $\square$

Proof of Theorem 4.5. Let $l = (0, \dots, 0, 1) \in \mathbb{R}^{d+1}$. From our definition of ε we clearly have

$$|b_k^\alpha - \hat{b}_k^\alpha| + m^\alpha |f^\alpha - \hat{f}^\alpha| + |c^\alpha - \hat{c}^\alpha| \leq \varepsilon.$$

Assume initially that $\varepsilon \in (0, h]$. Then, using that for any $a, b \geq 0$,

$$|a^2 - b^2| = (a + b)|a - b| = 2(a \wedge b)|a - b| + |a - b|^2$$

we get

$$|a_k^\alpha - \hat{a}_k^\alpha| \leq (|\sigma_k^\alpha| \wedge |\hat{\sigma}_k^\alpha|) |\sigma_k^\alpha - \hat{\sigma}_k^\alpha| + |\sigma_k^\alpha - \hat{\sigma}_k^\alpha|^2 \leq 2\varepsilon \sqrt{a_k^\alpha \wedge \hat{a}_k^\alpha} + \varepsilon h.$$

So we can use 4.6 for $Q = \mathcal{M}_T$ to get

$$\begin{aligned} & |u - \hat{u}| \\ & \leq \varepsilon N e^{c_0 T'} \left[1 + |u|_{0, Q} + |\hat{u}|_{0, Q} + \sup_{k, \partial Q} (|\delta_{h_k, \ell_k} u| + |\delta_{h_k, \ell_k} \hat{u}| + \varepsilon^{-1}|u - \hat{u}|) \right], \end{aligned}$$

on $\mathcal{M}_T$. Use Corollary 3.5 to estimate the term $|u|_{0, \mathcal{M}_T} + |\hat{u}|_{0, \mathcal{M}_T}$ to obtain

$$|u - \hat{u}| \leq \varepsilon N T e^{c_0(T+\tau)} I.$$

Drop the initial assumption that $\varepsilon \leq h$. For $\theta \in [0, 1]$, let u^θ be the solution of

$$\begin{aligned} \sup_\alpha m^\alpha (\delta_\tau u^\theta + a_k^{\theta\alpha} \Delta_{h_k, \ell_k} u^\theta + b_k^{\theta\alpha} \delta_{h_k, \ell_k} u^\theta - c^{\theta\alpha} u^\theta + f^{\theta\alpha}) &= 0 \quad \text{on } \mathcal{M}_T \\ (1 - \theta)u + \theta \hat{u} &= u^\theta \quad \text{on } \{(T, x) \in \bar{\mathcal{M}}_T\}, \end{aligned}$$

where

$$(\sigma_k^{\theta\alpha}, b_k^{\theta\alpha}, c^{\theta\alpha}, f^{\theta\alpha}) = (1 - \theta)(\sigma_k^\alpha, b_k^\alpha, c^\alpha, f^\alpha) + \theta(\hat{\sigma}_k^\alpha, \hat{b}_k^\alpha, \hat{c}^\alpha, \hat{f}^\alpha)$$

and $a_k^{\theta\alpha} = (1/2)|\sigma_k^{\theta\alpha}|^2$. For any $\theta_1, \theta_2 \in [0, 1]$,

$$|\sigma_k^{\theta_1\alpha} - \sigma_k^{\theta_2\alpha}| + |b_k^{\theta_1\alpha} - b_k^{\theta_2\alpha}| + |c^{\theta_1\alpha} - c^{\theta_2\alpha}| + m^\alpha |f^{\theta_1\alpha} - f^{\theta_2\alpha}| \leq |\theta_1 - \theta_2| \varepsilon.$$

Hence if θ_1, θ_2 satisfy $|\theta_1 - \theta_2| \varepsilon \leq h$, then, thanks to the first part of the proof (where u^{θ_1} would play the part of u while u^{θ_2} would play the part of $\hat{u}$),

$$|u^{\theta_1} - u^{\theta_2}| \leq N |\theta_1 - \theta_2| \varepsilon T' e^{c_0(T+\tau)} I(\theta_1, \theta_2),$$

with

$$\begin{aligned} I(\theta_1, \theta_2) &:= \sup_{\partial_\varepsilon Q} \left(1 + \max_k (|\delta_{h_k, \ell_k} u^{\theta_1}| + |\delta_{h_k, \ell_k} u^{\theta_2}|) \right. \\ & \quad \left. (|\theta_1 - \theta_2| \varepsilon)^{-1} |u^{\theta_1}(T, x) - u^{\theta_2}(T, x)| \right) \leq 4I, \end{aligned}$$

since $u^\theta = (1 - \theta)u + \theta\hat{u}$ on $\partial_\varepsilon Q = \{(T, x) \in \bar{\mathcal{M}}_T\}$. Subdivide the interval $[0, 1]$ into intervals of appropriate length to complete the proof for any $\varepsilon > 0$. $\square$

5. SOME PROPERTIES OF PAYOFF FUNCTIONS

We assume in the whole section that Assumption 2.1 and the following assumption hold.

Assumption 5.1. There is a function $m : A \rightarrow (0, 1]$, such that for all $\alpha \in A$, $t \in [0, T)$, $x \in \mathbb{R}^d$ and $y \in \mathbb{R}^d$

$$m^\alpha c^\alpha(t, x) + m^\alpha \geq K^{-1}, \quad |m^\alpha f^\alpha(t, x)| \leq K, \quad (5.1)$$

$$m^\alpha |f^\alpha(t, x) - f^\alpha(t, y)| \leq K|x - y|. \quad (5.2)$$

Lemma 5.2. *There exists a constant N such that:*

- (1) For any $(s, x) \in [0, T] \times \mathbb{R}^d$,

$$v(s, x) \leq N(1 + (T - s)).$$

- (2) For any $(s, x), (s, y) \in [0, T] \times \mathbb{R}^d$,

$$|v(s, x) - v(s, y)| \leq N(T - s)^2 e^{N(T-s)} |x - y|.$$

Proof. Since $m^\alpha c^\alpha(t, x) + m^\alpha \geq K^{-1}$ and $|m^\alpha f^\alpha(s, x)| \leq K$,

$$|f^\alpha(s, x)| \leq K^2(1 + c^\alpha(s, x)).$$

This and the assumption on the polynomial growth of g together with the estimates of moments for solutions to SDEs gives

$$\begin{aligned} |v^\alpha(s, x)| &\leq \mathbb{E}_{s,x}^\alpha \int_0^{T-s} |f^{\alpha t}(s+t, x_t)| e^{-\varphi t} dt + \mathbb{E}_{s,x}^\alpha |g(x_{T-s})| e^{-\varphi(T-s)} \\ &\leq K^2 \mathbb{E}_{s,x}^\alpha \int_0^{T-s} (1 + c^{\alpha t}(s+t, x_t)) e^{-\int_0^t c^{\alpha u}(s+u, x_u) du} dt + N \\ &\leq N(1 + (T - s)) + K^2 \mathbb{E} \int_0^{T-s} c^{\alpha t}(s+t, x_t) e^{-\int_0^t c^{\alpha u}(s+u, x_u) du} dt \\ &\leq N(1 + (T - s)). \end{aligned}$$

The right hand side of the estimate is independent of α , hence taking the supremum completes the first part of the proof. If $x_t^{s,x}$ is a solution to a stochastic differential equation

$$x_t^{s,x} = x + \int_0^t \sigma(s+u, x_u) dw_u + \int_0^t \beta(s+u, x_u) du$$

where

$$|\sigma(t, x) - \sigma(t, y)| \leq L|x - y| \quad \text{and} \quad |\beta(t, x) - \beta(t, y)| \leq L|x - y|,$$

then, by Theorem 2.5.9 from [5], with N depending on L only,

$$\mathbb{E} \sup_{t \leq T-s} |x_t^{s,x} - x_t^{s,y}| \leq N e^{N(T-s)} |x - y|. \quad (5.3)$$

First get an estimate for a fixed $\alpha \in A$.

$$\begin{aligned} |v^\alpha(s, x) - v^\alpha(s, y)| &\leq \mathbb{E} \int_0^{T-s} |f^{\alpha_t}(s+t, x_t^{s,x})| |e^{-\varphi_t^{\alpha,s,x}} - e^{-\varphi_t^{\alpha,s,y}}| dt \\ &\quad + \mathbb{E} \int_0^{T-s} |f^{\alpha_t}(s+t, x_t^{s,x}) - f^{\alpha_t}(s+t, x_t^{s,y})| e^{-\varphi_t^{\alpha,s,y}} dt \\ &\quad + \mathbb{E} |g(x_{T-s}^{s,x}) - g(x_{T-s}^{s,y})| =: I_1 + I_2 + I_3. \end{aligned}$$

Estimating the above integrals separately:

$$I_1 \leq \mathbb{E} \int_0^{T-s} |f^{\alpha_t}(s+t, x_t^{s,x})| |\varphi_t^{\alpha,s,x} - \varphi_t^{\alpha,s,y}| e^{-\min(\varphi_t^{\alpha,s,x}, \varphi_t^{\alpha,s,y})} dt.$$

Since $m^\alpha c^\alpha(t, x) + m^\alpha \geq K^{-1}$ and $|m^\alpha f^\alpha(s, x)| \leq K$,

$$|f^\alpha(s, x)| \leq \frac{K}{m^\alpha} \quad \text{and} \quad c^\alpha(s, x) \geq \frac{1}{Km^\alpha} - 1.$$

Therefore, using Lipschitz continuity of c when estimating $|\varphi_t^{\alpha,s,x} - \varphi_t^{\alpha,s,y}|$,

$$\begin{aligned} I_1 &\leq (T-s) e^{T-s} K^3 \mathbb{E} \int_0^{T-s} \frac{1}{Km^{\alpha_t}} \sup_{r \leq t} |x_r^{\alpha,s,x} - x_r^{\alpha,s,y}| e^{-\int_0^t \frac{1}{Km^{\alpha_u}} du} dt \\ &\leq (T-s) e^{T-s} K^3 \mathbb{E} \sup_{t \leq T-s} |x_t^{\alpha,s,x} - x_t^{\alpha,s,y}| \int_0^{T-s} -de^{-\int_0^t \frac{1}{Km^{\alpha_u}} du}. \end{aligned}$$

Now

$$\begin{aligned} I_2 &\leq \int_0^{T-s} (m^{\alpha_t})^{-1} |x_t^{\alpha,s,x} - x_t^{\alpha,s,y}| e^{-\varphi_t^{\alpha,s,y}} dt \\ &\leq \mathbb{E} \left(\sup_{t \leq T-s} |x_t^{\alpha,s,x} - x_t^{\alpha,s,y}| \int_0^{T-s} K(1 + c^{\alpha_t}(s+t, x_t^{\alpha,s,y})) e^{-\varphi_t^{\alpha,s,y}} dt \right) \\ &\leq [K(t-s) + K] \mathbb{E} \sup_{t \leq T-s} |x_t^{\alpha,s,x} - x_t^{\alpha,s,y}|. \end{aligned}$$

and

$$I_3 \leq K(T-s) \mathbb{E} \sup_{t \leq T-s} |x_t^{\alpha,s,x} - x_t^{\alpha,s,y}|.$$

Hence

$$|v^\alpha(s, x) - v^\alpha(s, y)| \leq K^3(T-s)^2 e^{(T-s)} \mathbb{E} \sup_{t \leq T-s} |x_t^{\alpha,s,x} - x_t^{\alpha,s,y}|.$$

By (5.3) the right hand side is independent of α . $\square$

Let $H_T := [0, T) \times \mathbb{R}^d$ and $\partial H_T := \{T\} \times \mathbb{R}^d$.

Lemma 5.3. *Let ψ be a smooth function on H_T such that its first order partial derivatives in x grow at most polynomially and for all $\alpha \in A$ it satisfies*

$$\frac{\partial}{\partial t} \psi + L^\alpha \psi - c^\alpha \psi + f^\alpha \leq 0 \quad \text{on } H_T.$$

Let v be the payoff function of the stochastic control problem (2.5). Then

$$v \leq \psi + \sup_{\partial H_T} [v - \psi]_+ \quad \text{on } H_T. \quad (5.4)$$

Hence, clearly, if $v \leq \psi$ on ∂H_T , then $v \leq \psi$ on H_T .

Proof. For any $\varepsilon > 0$ there is a control process $\alpha_t \in \mathfrak{A}$ such that

$$\begin{aligned} v(s, x) &\leq \varepsilon + \mathbb{E}_{s,x} \left(\int_0^{T-s} f^{\alpha_t}(s+t, x_t) e^{-\varphi_t} dt + g(x_{T-s}) e^{-\varphi_{T-s}} \right) \\ &\leq \varepsilon - \mathbb{E}_{s,x} \int_0^{T-s} e^{-\varphi_t} \left[\frac{\partial \psi}{\partial t}(s+t, x_t) + L^{\alpha_t} \psi(s+t, x_t) - c^{\alpha_t} \psi(s+t, x_t) \right] dt \\ &\quad + \mathbb{E}_{s,x} g(x_{T-s}) e^{-\varphi_{T-s}}. \end{aligned}$$

Applying Itô's formula to $\psi(s+r, x_r) e^{-\varphi_r}$ on the interval $[0, T-s]$ and taking expectation we get

$$\begin{aligned} \psi(s, x) &= \mathbb{E}_{s,x} \psi(T, x_{T-s}) e^{-\varphi_{T-s}} \\ &\quad - \mathbb{E}_{s,x} \int_0^{T-s} e^{-\varphi_t} \left[\frac{\partial}{\partial t} + L^\alpha - c^\alpha \right] \psi(s+t, x_t) dt, \end{aligned}$$

by noting that the Itô integral has zero expectation, due to moment estimates for x_r , since ψ has polynomially growing first order partial derivatives in x . Hence

$$v(s, x) \leq \varepsilon + \psi(t, x) + \sup_{x \in \mathbb{R}^d} [v(T, x) - \psi(T, x)]_+,$$

for any $\varepsilon > 0$, which yields (5.4). $\square$

Lemma 5.4. *Let ψ be a smooth function on H_T such that its first order partial derivatives in x grow at most polynomially and there exists $\alpha_0 \in A$ such that*

$$\frac{\partial}{\partial t} \psi + L^{\alpha_0} \psi - c^{\alpha_0} \psi + f^{\alpha_0} \geq 0 \quad \text{on } H_T.$$

Let v be the payoff function of the stochastic control problem (2.5). Then

$$\psi \leq v + \sup_{\partial H_T} [\psi - v]_+ \quad \text{on } H_T.$$

Hence, clearly, if $\psi \leq v$ on ∂H_T , then $\psi \leq v$ on H_T .

Proof. For the constant strategy $\alpha_t := \alpha_0$ for all $t \in [0, T-s]$

$$\begin{aligned} v(s, x) &\geq \mathbb{E}_{s,x}^\alpha \left(\int_0^{T-s} f^{\alpha_0}(s+t, x_t) e^{-\varphi_t} dt + g(x_{T-s}) e^{-\varphi_{T-s}} \right) \\ &\geq \mathbb{E}_{s,x}^\alpha \left(\int_0^{T-s} - \left[\frac{\partial}{\partial t} + L^{\alpha_0} - c^{\alpha_0} \right] \psi(s+t, x_t) e^{-\varphi_t} dt + g(x_{T-s}) e^{-\varphi_{T-s}} \right). \end{aligned}$$

By Itô's formula for $\psi(s+r, x_r) e^{-\varphi_r}$ on the interval $[0, T-s]$

$$\psi(s, x) \leq v(s, x) + \sup_{x \in \mathbb{R}^d} [\psi(T, x) - v(T, x)]_+.$$

$\square$

Corollary 5.5. *Assume (2.11) holds and for all $\alpha \in A$ and $(t, x) \in H_T$*

$$m^\alpha |f^\alpha| \leq K \quad \text{and} \quad m^\alpha c^\alpha \leq K. \quad (5.5)$$

Then there exists a constant N such that for $(s_0, x_0) \in H_T$ and $(t_0, x_0) \in H_T$ satisfying $s_0 - 1 \leq t_0 \leq s_0$ one has

$$|v(s_0, x_0) - v(t_0, x_0)| \leq N(\nu + 1) |s_0 - t_0|^{\frac{1}{2}},$$

where ν is the Lipschitz constant from part 2 of Lemma 5.2.

Proof. Shift the time axis so that $t_0 = 0$, $s_0 \leq 1$. Let $(t, x) \in H_{s_0} := [0, s_0] \times \mathbb{R}^d$. Define $\xi(t) = e^{s_0-t}$. Let $\zeta = \xi\eta$, $\eta(x) = |x - x_0|^2$ and $\gamma := s_0^{-1/2}$. Let

$$\psi(t, x) = \gamma\nu [\zeta(t, x) + \kappa(s_0 - t)] + \frac{1}{\gamma}\nu\kappa + v(s_0, x_0),$$

where $\kappa > 1$ is a large positive constant to be chosen later. To apply Lemma 5.3 we need to show that for all $\alpha \in A$

$$m^\alpha \left(\frac{\partial}{\partial t} \psi + L^\alpha \psi - c^\alpha \psi + f^\alpha \right) \leq 0 \quad \text{on } H_{s_0}$$

and $\psi(s_0, x) \geq v(s_0, x)$ for any $x \in \mathbb{R}^d$. Notice

$$\frac{\partial}{\partial t} \psi = -\gamma\nu\eta\xi - \gamma\nu\kappa, \quad \frac{\partial}{\partial x_i} \eta(x) = 2(x^i - x_0^i), \quad \frac{\partial^2}{\partial x_i \partial x_j} \eta = 2\delta_{i,j},$$

where $\delta_{i,j}$ is the Kronecker delta. Using (2.11) and $\xi \leq e^T$,

$$L^\alpha \psi \leq \gamma\nu 2Kd(1 + |x - x_0|)\xi \leq \gamma\nu 2Kd(1 + |x - x_0|)e^T.$$

Since v is bounded by N_T , for any $\alpha \in A$ and $(t, x) \in H_{s_0}$

$$-c^\alpha(t, x)v(s_0, x_0) \leq c^\alpha(t, x)N_T.$$

Hence for any $\alpha \in A$

$$-c^\alpha \psi \leq -c^\alpha \frac{1}{\gamma} \nu \kappa + c^\alpha N_T \quad \text{on } H_{s_0}.$$

Then for any $\alpha \in A$ and $(t, x) \in H_{s_0}$, with $N_2 := 2Ke^T d$ and by (5.5),

$$\begin{aligned} & m^\alpha \left(\frac{\partial}{\partial t} \psi + L^\alpha \psi - c^\alpha \psi + f^\alpha \right) \\ & \leq m^\alpha \left\{ \gamma\nu [-|x - x_0|^2 + N_2(1 + |x - x_0|) - \kappa] - c^\alpha \frac{\kappa}{\gamma} \nu + c^\alpha N_T \right\} + K. \end{aligned}$$

The right hand side is a quadratic in $|x - x_0|$ with a negative leading coefficient and so it achieves its maximum when $|x - x_0| = N_2/2$. The maximum is

$$m^\alpha \left\{ \gamma\nu \left[-\frac{N_2}{4} + N_2 + \frac{N_2^2}{2} - \kappa \right] + c^\alpha \kappa \gamma^{-1} \nu + c^\alpha N_T \right\} + K \leq M - L\kappa, \quad (5.6)$$

where $L := K^{-1}\nu$ and M depends only on N , d and T . Indeed, $s_0 \leq 1$ so $\gamma \geq 1$. Since $m^\alpha(1 + c^\alpha) \geq K^{-1}$ on H_T , $m^\alpha(\gamma^2 + c^\alpha) \geq (1 \wedge \gamma^2)K^{-1}$ on H_T . Hence

$$\kappa\nu\gamma^{-1}m^\alpha(\gamma^2 + c^\alpha) \geq \kappa\nu\gamma^{-1}(\gamma^2 \wedge 1)K^{-1} \geq K^{-1}\nu\kappa = L\kappa.$$

If κ is large (depending on K, T, d only) then the right hand side of (5.6) is non-positive. Hence

$$m^\alpha \left(\frac{\partial}{\partial t} \psi + L^\alpha \psi - c^\alpha \psi + f^\alpha \right) \leq 0.$$

By Lemma 5.2

$$\begin{aligned} \psi(s_0, x) &= \gamma\nu|x - x_0|^2 + \nu\kappa\gamma^{-1} + v(s_0, x_0) \\ &\geq \gamma\nu|x - x_0|^2 + \nu\kappa\gamma^{-1} - \nu|x - x_0| + v(s_0, x). \end{aligned}$$

The reader can check that

$$\gamma\nu|x-x_0|^2 + \nu\kappa\gamma^{-1} - \nu|x-x_0| \geq 0,$$

since it is a quadratic in $|x-x_0|$, its leading coefficient is positive and hence its minimum is $4^{-1}\nu\gamma^{-1} + \nu\kappa\gamma^{-1} - 2^{-1}\nu\gamma^{-1}$. Hence for all $x \in \mathbb{R}^d$,

$$\psi(s_0, x) \geq v(s_0, x).$$

By Lemma 5.3, $\psi(t_0, x_0) \geq v(t_0, x_0)$. But

$$v(t_0, x_0) \leq \psi(t_0, x_0) = \gamma\nu\kappa s_0 + \gamma^{-1}\nu\kappa + v(s_0, x_0).$$

Hence, recalling that $\gamma = s_0^{-1/2}$,

$$v(t_0, x_0) \leq v(s_0, x_0) + s_0^{-1/2}\nu\kappa s_0 + s_0^{1/2}\nu\kappa \leq v(s_0, x_0) + N_T(\nu + 1)s_0^{1/2}.$$

What remains to be shown is that

$$v(s_0, x_0) \leq v(t_0, x_0) + N_T(\nu + 1)s_0^{1/2}.$$

But that can be achieved by considering

$$\psi = -\gamma\nu[\zeta + \kappa(s_0 - t)] - \frac{1}{\gamma}\nu\kappa + v(s_0, x_0)$$

and applying Lemma 5.4 to ψ . $\square$

6. SHAKING THE COEFFICIENTS

The method of shaking the coefficients first introduced in [7] and [8] will be used. Recall that $v_{\tau,h}$ is the solution of (2.9)-(2.10) with $Q = \mathcal{M}_T$. Recall that v is the payoff function (2.5).

Remark 6.1. For (t_0, x_0) not in $\bar{\mathcal{M}}_T$ define $v_{\tau,h}(t_0, x_0)$ as the solution to (2.9)-(2.10) on

$$\begin{aligned} \mathcal{M}_T(t_0, x_0) = \{ & ((t_0 + j\tau) \wedge T, x_0 + h(i_1\ell_1 + \dots + i_{d_1}\ell_{d_1})) : \\ & j \in \mathbb{N} \cup \{0\}, i_k \in \mathbb{Z}, k = 1, \dots, d_1 \}. \end{aligned}$$

Let B_1 denote the unit ball centered at the origin, in R^d . Consider $y \in S \subset B_1$ and $r \in \Lambda \subset (-1, 0)$. Fix $\varepsilon > 0$. Let $v_{\tau,h}^\varepsilon$ be the unique solution of

$$\begin{aligned} \sup_{\alpha \in A, y \in S, r \in \Lambda} m^\alpha \Big[& \delta_\tau u(t, x) + L_h^\alpha(t + \varepsilon^2 r, x + \varepsilon y)u(t, x) \\ & + f^\alpha(t + \varepsilon^2 r, x + \varepsilon y) \Big] = 0 \quad \text{on } \mathcal{M}_T, \\ u(T, x) = & \sup_{y \in S} g(x + \varepsilon y). \end{aligned} \tag{6.1}$$

Assumption 2.4 is satisfied by (6.1) and so the solution exists, is unique and has all the other properties proved in section 3. Assume, from now on, that τ is small enough, so that by Remark 4.4, (4.10) holds.

Lemma 6.2. *Let Assumption 2.4 hold. Then there is a constant N , such that*

$$|v_{\tau,h}^\varepsilon - v_{\tau,h}| \leq Ne^{NT}T\varepsilon.$$

Proof. Let the space of controls be $A \times S \times \Lambda$. Let

$$(\hat{\sigma}_k^\alpha, \hat{b}_k^\alpha, \hat{c}^\alpha, \hat{f}^\alpha)(t, x) := (\sigma_k^\alpha, b_k^\alpha, c^\alpha, f^\alpha)(t + \varepsilon^2 r, x + \varepsilon y).$$

Then

$$\sup_{\mathcal{M}_T, A \times S \times \Lambda, k} \left(|\sigma_k^\alpha - \hat{\sigma}_k^\alpha| + |b_k^\alpha - \hat{b}_k^\alpha| + |c^\alpha - \hat{c}^\alpha| + m^\alpha |f^\alpha - \hat{f}^\alpha| \right) \leq 8K\varepsilon.$$

Apply Theorem 4.5 to get the conclusion. $\square$

The Lipschitz continuity of the solutions to (2.9)-(2.10) follows. Consider a particular case of (6.1). Let $\Lambda = \{0\}$. Let $\varepsilon = |x - y|$. Fix $x, y \in \mathbb{R}^d$ and let $S = \{\varepsilon^{-1}(x - y)\}$. Say $z \in S$. Then by uniqueness

$$v_{\tau, h}^\varepsilon(t, x) = v_{\tau, h}(x - \varepsilon z) = v_{\tau, h}(t, y).$$

Hence one obtains:

Corollary 6.3. *Let Assumption 2.4 hold. Then there is a constant N such that for any $t \in [0, T]$ and $x, y \in \mathbb{R}^d$,*

$$|v_{\tau, h}(t, x) - v_{\tau, h}(t, y)| \leq NTe^{NT}|x - y|. \quad (6.2)$$

Corollary 6.4. *Let Assumptions 2.4 and 2.5 hold. Then there is a constant N such that for any $s, t \in [0, T]$ and $x \in \mathbb{R}^d$,*

$$|v_{\tau, h}(t, x) - v_{\tau, h}(s, x)| \leq NTe^{NT}|t - s|^{1/2}. \quad (6.3)$$

The proof of this corollary is identical to the proof of (6.8) of Lemma 6.2 in [1], if one uses Lemma 3.7, Corollary 6.3 and Theorem 4.5 instead of Corollary 3.7, Lemma 6.1 and Theorem 5.6 from [1], respectively.

In the continuous case, the shaking will be introduced as follows. Let Assumptions 2.1 and 5.1 hold. Consider the separable metric space

$$C = A \times \{(\tau, \xi) \in (-1, 0) \times B_1\},$$

with the metric which comes from taking the sum of the metric for A and the metrics which are induced by natural norms on $(-1, 0)$ and B_1 . Extend all the functions σ, β, f, c for negative t by $\sigma^\gamma(t, x) = \sigma^\gamma(0, x)$ etc. For a fixed $\varepsilon \in (0, 1)$, for $\gamma = (\alpha, \tau, \xi)$ let

$$\sigma^\gamma(t, x) = \sigma^\alpha(t + \varepsilon^2 \tau, x + \varepsilon \xi) \quad (6.4)$$

and similarly for β, c and f . Let $x_t^{\gamma, s, x}$ be the solution of

$$x_t^{s, x} = x + \int_0^t \sigma^\gamma(s + u, x_u) dw_u + \int_0^t \beta^\gamma(s + u, x_u) du. \quad (6.5)$$

Let $C_n := A_n \times \{(\tau, \xi) \in (-1, 0) \times B_1\}$, $C = \bigcup_{n \in \mathbb{N}} C_n$. Let $\mathfrak{C}_n$ be the spaces of admissible control processes defined analogously to $\mathfrak{A}_n$ and $\mathfrak{C}$ defined analogously to $\mathfrak{A}$. Let

$$w^\gamma(s, x) = \mathbb{E}_{s, x}^\gamma \left[\int_0^{T-s} f^\gamma(t, x_t) e^{-\varphi t} dt + g(x_{T-s}) e^{-\varphi(T-s)} \right], \quad (6.6)$$

$$w_n(s, x) = \sup_{\gamma \in \mathfrak{C}_n} w^\gamma(s, x), \quad (6.7)$$

$$w(s, x) = \sup_{\gamma \in \mathfrak{C}} w^\gamma(s, x). \quad (6.8)$$

Lemma 6.5. *If w^γ and w are defined by (6.6) and (6.8), respectively, then*

$$|w^\gamma - v^\alpha| \leq Ne^{NT}\varepsilon \quad \text{and} \quad |w - v| \leq Ne^{NT}\varepsilon \quad \text{on } H_T. \quad (6.9)$$

Proof. Let $x^{\gamma,s,x}$ and $x^{\alpha,s,x}$ denote the solutions to (6.5) and (2.2) respectively. By Theorem 2.5.9 of [5]

$$\begin{aligned} \mathbb{E} \sup_{t \leq T-s} |x_t^{\gamma,s,x} - x_t^{\alpha,s,x}|^2 &\leq Ne^{NT} \sup [|\sigma^\alpha(t + \varepsilon^2\tau, x + \varepsilon\xi) - \sigma^\alpha(t, x)|^2 \\ &\quad + |\beta^\alpha(t + \varepsilon^2\tau, x + \varepsilon\xi) - \beta^\alpha(t, x)|^2]. \end{aligned} \quad (6.10)$$

Hence, using Hölder inequality

$$\mathbb{E} \sup_{t \leq T-s} |x_t^{\gamma,s,x} - x_t^{\alpha,s,x}| \leq Ne^{NT}\varepsilon. \quad (6.11)$$

With this in mind the reader could use the same technique used in proving the second part of Lemma 5.2 in order to get the desired estimate. $\square$

Theorem 6.6. *For any $\varepsilon \in (0, 1]$ there exists u in $C^\infty([0, T] \times \mathbb{R}^d)$ such that*

$$\sup_{\alpha \in A} (u_t + L^\alpha u + f^\alpha) \leq 0 \quad \text{on } H_{T-\varepsilon^2} \quad (6.12)$$

$$|u - v| \leq Ne^{NT}\varepsilon \quad \text{on } H_T, \quad (6.13)$$

$$|D_t^2 u|_{0,[0,T] \times \mathbb{R}^d} + |D_x^4 u|_{0,[0,T] \times \mathbb{R}^d} \leq Ne^{NT}\varepsilon^{-3} \quad \text{on } H_T, \quad (6.14)$$

$$|D_x^2 u|_{0,[0,T] \times \mathbb{R}^d} \leq Ne^{NT}\varepsilon^{-1} \quad \text{on } H_T.$$

Proof. Recall that $A = \bigcup_{n \in \mathbb{N}} A_n$. By Theorem 2.1 of [8], there exist a smooth function u_n defined on H_T such that for all $\alpha \in A_n$,

$$\frac{\partial}{\partial t} u_n + L^\alpha u_n + f^\alpha \leq 0 \quad \text{on } H_{T-\varepsilon^2}. \quad (6.15)$$

Let w_n be defined by (6.7). Notice that f^γ , c^γ , σ^γ and b^γ in the definition all depend on ε and x_t is defined by (6.5). Let $\zeta \in C_0^\infty((-1, 0) \times B_1)$ be non-negative with unit integral. Let $\zeta_\varepsilon(t, x) := \varepsilon^{-d-2}\zeta(t/\varepsilon^2, x/\varepsilon)$. By the proof of Theorem 2.1 of [8], $u_n = w_n * \zeta_\varepsilon$. Let w be defined by (6.8) and let $u := w * \zeta_\varepsilon$. By Lemma (5.2) the functions w, w_n are bounded in absolute value by a constant independent of n . Then by Lebesgue Dominated Convergence Theorem

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\partial}{\partial t} u_n(t, x) &= \lim_{n \rightarrow \infty} \frac{\partial}{\partial t} \int_{(-1,0) \times B_1} \zeta_\varepsilon(t-s, x-y) w_n(s, y) ds dy \\ &= \int_{(-1,0) \times B_1} \lim_{n \rightarrow \infty} w_n(s, y) \frac{\partial}{\partial t} \zeta_\varepsilon(t-s, x-y) ds dy \\ &= \frac{\partial}{\partial t} u(t, x). \end{aligned}$$

Hence $\lim_{n \rightarrow \infty} \frac{\partial}{\partial t} u_n = \frac{\partial}{\partial t} u$. Similarly $\lim_{n \rightarrow \infty} L^\alpha u_n = L^\alpha u$ on H_T , for any $\alpha \in A$. For each fixed $\alpha \in A$, let $n \rightarrow \infty$ in (6.15). Then for any $\alpha \in A$

$$\frac{\partial}{\partial t} u + L^\alpha u + f^\alpha \leq 0 \quad \text{on } H_{T-\varepsilon^2},$$

which shows (6.12). By Lemma 5.2 and Corollary 5.5

$$|w(t, x) - w(s, y)| \leq Ne^{NT}(|t-s|^{1/2} + |x-y|).$$

Then (6.14) follows from known properties of convolutions. Now we want to show (6.13). Since $u = w * \zeta_\varepsilon$, $|u(s, x) - w(s, x)|$ is estimated by the right hand side of (6.13). So we need only estimate $|w(s, x) - v(s, x)|$, for $(s, x) \in H_T$. To this end we use the fact that difference of supremums is less than the supremum of a difference to see that we need only estimate

$$\begin{aligned} & \mathbb{E}_{s,x} \int_0^{T-s} |f^{\gamma_t}(s+t, x_t^\gamma) e^{-\varphi_t^\gamma} - f^{\alpha_t}(s+t, x_t^\alpha) e^{-\varphi_t^\alpha}| dt, \\ & + \mathbb{E}_{s,x} |g(x_{T-s}^\gamma) e^{-\varphi_{T-s}^\gamma} - g(x_{T-s}^\alpha) e^{-\varphi_{T-s}^\alpha}| =: I + J. \end{aligned}$$

Clearly

$$\begin{aligned} J & \leq K \left[T2\varepsilon + 2\mathbb{E}_{s,x} \sup_{t \leq T-s} |x_t^\gamma - x_t^\alpha| \right], \\ I & \leq \mathbb{E}_{s,x} \int_0^{T-s} |f^{\gamma_t}(s+t, x_t^\gamma)| |e^{-\varphi_t^\gamma} - e^{-\varphi_t^\alpha}| dt \\ & + \mathbb{E}_{s,x} \int_0^{T-s} |f^{\gamma_t}(s+t, x_t^\gamma) - f^{\alpha_t}(s+t, x_t^\alpha)| e^{-\varphi_t^\alpha} dt =: I_1 + I_2. \end{aligned}$$

Like in the proof of Lemma 5.2 (using Assumptions 2.4, 2.5 and the definition of c^γ), we get

$$\begin{aligned} I_1 & \leq \mathbb{E}_{s,x} \int_0^{T-s} K^2 (m^{\alpha_t})^{-1} t \left[2\varepsilon + \sup_{u \leq t} |x_u^\gamma - x_u^\alpha| \right] e^{-\min(\varphi_t^\gamma, \varphi_t^\alpha)} dt \\ & \leq K^3 e^T \left[2\varepsilon + \mathbb{E}_{s,x} \sup_{t \leq T-s} |x_t^\gamma - x_t^\alpha| \right]. \end{aligned}$$

By Assumption 2.4 and the definition of f^γ

$$\begin{aligned} I_2 & = \mathbb{E}_{s,x} \int_0^{T-s} |f^{\alpha_t}(s+t+\varepsilon^2\tau, x_t^\gamma + \varepsilon\xi) - f^{\alpha_t}(s+t, x_t^\alpha)| e^{-\varphi_t^\alpha} dt \\ & \leq \mathbb{E}_{s,x} \int_0^{T-s} |K(m^{\alpha_t})^{-1}| [2\varepsilon + |x_t^\gamma - x_t^\alpha|] e^{-\varphi_t^\alpha} dt \\ & \leq 2e^T K^2 \left[\varepsilon + \mathbb{E}_{s,x} \sup_{t \leq T-s} |x_t^\gamma - x_t^\alpha| \right]. \end{aligned}$$

By Theorem 2.5.9 of [5]

$$\mathbb{E}_{s,x} \sup_{u \leq T-s} |x_u^\gamma - x_u^\alpha| \leq N e^{NT} \varepsilon,$$

where x^α is a solution of (2.2), while x^γ is a solution of (6.5). Hence, noting that the estimate for $I + J$ is independent of α , τ and ξ , we get (6.13). $\square$

Lemma 6.7. *Let Assumptions 2.1, 2.3, 2.4 and 2.5 hold. Then*

$$v \leq v_{\tau,h} + N_T(\tau^{1/4} + h^{1/2}). \quad (6.16)$$

Proof. Recall that for $\varepsilon > 0$, $v_{\tau,h}^\varepsilon$ is defined as the unique bounded solution to (6.1) and by Lemma 6.2

$$v_{\tau,h}^\varepsilon \leq v_{\tau,h} + N\varepsilon.$$

Assume, without loss of generality, that $h \leq 1$. Let $\xi \geq 0$ be a $C_0^\infty([0, T] \times \mathbb{R}^d)$ function with a unit integral and support in $(-1, 0) \times B_1$. For any function w defined on $(-\infty, T) \times \mathbb{R}^d$, for which it makes sense, let

$$w^{(\varepsilon)}(t, x) = \int_{H_T} w(t - \varepsilon^2 r, x - \varepsilon y) \xi(r, y) dr dy.$$

If the function $w(t, x)$ is not defined for negative t then extend it for $t < 0$ by defining $w(t, x) = w(0, x)$. For any $\alpha \in A$ and for all $r \in \Lambda$, $y \in S$

$$\begin{aligned} \delta_\tau v_{\tau, h}^\varepsilon(t - \varepsilon^2 r, x - \varepsilon y) + L_h^\alpha(t, x) v_{\tau, h}^\varepsilon(t - \varepsilon^2 r, x - \varepsilon y) \\ + f^\alpha(t, x) \leq 0 \quad \text{on } H_T. \end{aligned}$$

Multiply this by $\xi(r, y)$ and integrate over H_T with respect to r and y . Then

$$\delta_\tau v_{\tau, h}^{\varepsilon(\varepsilon)} + L_h^\alpha v_{\tau, h}^{\varepsilon(\varepsilon)} + f^\alpha \leq 0 \quad \text{on } H_T.$$

Use Taylor's Theorem, to get that on $H_{T-2\varepsilon^2}$:

$$\begin{aligned} |\delta_\tau v_{\tau, h}^{\varepsilon(\varepsilon)} - D_t v_{\tau, h}^{\varepsilon(\varepsilon)}| &\leq N\tau |D_t^2 v_{\tau, h}^{\varepsilon(\varepsilon)}|_{0, H_{T-2\varepsilon^2}} =: M_1 \\ m^\alpha |L_h^\alpha v_{\tau, h}^{\varepsilon(\varepsilon)} - L^\alpha v_{\tau, h}^{\varepsilon(\varepsilon)}| &\leq Nh^2 |D_x^4 v_{\tau, h}^{\varepsilon(\varepsilon)}|_{0, H_{T-2\varepsilon^2}} \\ &\quad + h |D_x^2 v_{\tau, h}^{\varepsilon(\varepsilon)}|_{0, H_{T-2\varepsilon^2}} =: M_2. \end{aligned}$$

With $M := M_1 + M_2$, for any $\alpha \in A$ on $H_{T-2\varepsilon^2}$

$$[D_t + L^\alpha](v_{\tau, h}^{\varepsilon(\varepsilon)} + (T - t)M) + f^\alpha \leq 0.$$

By Lemma 5.3

$$v \leq v_{\tau, h}^{\varepsilon(\varepsilon)} + 2(T - t)M + \sup_{\partial H_{T-2\varepsilon^2}} (v - v_{\tau, h}^{\varepsilon(\varepsilon)})_+.$$

By Hölder continuity in time of v and also $v_{\tau, h}^{\varepsilon(\varepsilon)}$:

$$\begin{aligned} \sup_{\partial H_{T-2\varepsilon^2}} (v - v_{\tau, h}^{\varepsilon(\varepsilon)})_+ &\leq \sup_{x \in \mathbb{R}^d} (|v(T - \varepsilon^2, x) - g(x)| + |v_{\tau, h}^{\varepsilon(\varepsilon)}(T - \varepsilon^2, x) - g(x)|) \\ &\leq 4N_T \varepsilon. \end{aligned}$$

By standard properties of convolutions:

$$\begin{aligned} |D_t^2 v_{\tau, h}^{\varepsilon(\varepsilon)}|_{0, H_{T-2\varepsilon^2}} + |D_x^4 v_{\tau, h}^{\varepsilon(\varepsilon)}|_{0, H_{T-2\varepsilon^2}} + |D_x^2 v_{\tau, h}^{\varepsilon(\varepsilon)}|_{0, H_{T-2\varepsilon^2}} \\ \leq N\varepsilon^{-3} + N\varepsilon^{-3} + N\varepsilon^{-1}. \end{aligned}$$

Since $\varepsilon = (\tau + h^2)^{1/4}$,

$$v \leq v_{\tau, h}^{\varepsilon(\varepsilon)} + 4N\varepsilon + N\tau\varepsilon^{-3} + Nh\varepsilon^{-1} + Nh^2\varepsilon^{-3} \leq v_{\tau, h}^{\varepsilon(\varepsilon)} + N(\tau + h^2)^{1/4}.$$

Then

$$v \leq v_{\tau, h} + N(\tau^{1/4} + h^2) \quad \text{on } H_T,$$

since $|v_{\tau, h}^\varepsilon - v_{\tau, h}| \leq N\varepsilon$ and $|v_{\tau, h}^{\varepsilon(\varepsilon)} - v_{\tau, h}^\varepsilon| \leq N\varepsilon$. □

Lemma 6.8. *Let Assumptions 2.1, 2.3, 2.4 and 2.5 hold. Then*

$$v_{\tau, h} \leq v + N_T(\tau^{1/4} + h^{1/2}) \quad \text{on } H_T. \quad (6.17)$$

Proof. Let $\varepsilon = (\tau + h^2)^{1/4}$. On $(T - \varepsilon^2, T]$ the estimate is trivial consequence of the Hölder continuity in time of both v and $v_{\tau,h}$. Let $S := T - \varepsilon^2$. It remains to prove the estimate on H_S . By Theorem 6.6, there is a smooth function u defined on $[0, T] \times \mathbb{R}^d$ satisfying

$$\sup_{\alpha \in A} (u_t + L_h^\alpha u + f^\alpha) \leq 0 \quad \text{on } H_S. \quad (6.18)$$

Apply Taylor's Theorem to see that in $H_{T-\tau}$

$$\begin{aligned} |\delta_\tau u - D_t u| &\leq N\tau |D_t^2 u|_{0, H_T} =: M_1 \\ m^\alpha |L_h^\alpha u - L^\alpha u| &\leq Nh^2 |D_x^4 u|_{0, H_T} + h |D_x^2 u|_{0, H_T} =: M_2 \end{aligned} \quad (6.19)$$

Let $M = M_1 + M_2$. By Theorem 6.6:

$$M_1 + M_2 \leq N_T \tau \varepsilon^{-3} + N_T h^2 \varepsilon^{-3} + N_T h \varepsilon^{-1} \leq N_T (\tau + h^2)^{1/4}.$$

Since $\tau < 1$, $\varepsilon^2 > \tau$. By (6.18) and (6.19)

$$\begin{aligned} \sup_\alpha m^\alpha (\delta_\tau v_{\tau,h} + L_h^\alpha v_{\tau,h} + f^\alpha) &= 0 \\ &\geq \sup_\alpha (\delta_\tau u + L_h^\alpha u + f^\alpha - M) \quad \text{on } H_S. \end{aligned}$$

Let $u' = \sup_{H_T \setminus H_S} (v_{\tau,h} - u)_+ + u$. The aim now is to apply Lemma 3.3 to $v_{\tau,h}$ and u' . On the discrete boundary $H_T \setminus H_S$, $u' \geq v_{\tau,h}$. By Lemma 3.3

$$v_{\tau,h} \leq u + TM + \sup_{H_T \setminus H_S} (v_{\tau,h} - u)_+. \quad (6.20)$$

By Hölder continuity in time of $v_{\tau,h}$ and v and by Theorem 6.6

$$\sup_{H_T \setminus H_S} (v_{\tau,h} - u)_+ \leq \sup_{H_T \setminus H_S} |v_{\tau,h} - g| + \sup_{H_T \setminus H_S} |g - v| + \sup_{H_T \setminus H_S} |v - u| \leq N_T \varepsilon.$$

By Theorem 6.6, $|v - u| \leq N_T \varepsilon$. Hence

$$\begin{aligned} v_{\tau,h} &\leq v + N_T (\varepsilon + \tau^{1/2} + (\tau + h^2)^{1/4}) \leq v + N_T (\tau + h^2)^{1/4} \\ &\leq v + N_T (\tau^{1/4} + h^{1/2}) \quad \text{on } H_S. \end{aligned}$$

□

Acknowledgment. The authors are grateful to Nicolai Krylov in Minnesota for useful information on the subject of this paper.

REFERENCES

- [1] N. V. Krylov, "On the rate of convergence of finite-difference approximations for Bellman equations with Lipschitz coefficients," *arXiv:math*, vol. 1, no. 1, pp. 1–33, 2004.
- [2] A. N. Shiryaev, *Statisticheskii posledovatelnyi analiz. Optimalnye pravila ostanovki*. Izdat. "Nauka", Moscow, 1976. Second edition, revised.
- [3] J.-L. Menaldi, "Some estimates for finite difference approximations," *SIAM J. Control Optim.*, vol. 27, no. 3, pp. 579–607, 1989.
- [4] H. J. Kushner and P. Dupuis, *Numerical methods for stochastic control problems in continuous time*, vol. 24 of *Applications of Mathematics (New York)*. New York: Springer-Verlag, second ed., 2001. Stochastic Modelling and Applied Probability.
- [5] N. V. Krylov, *Controlled diffusion processes*, vol. 14 of *Applications of Mathematics*. New York: Springer-Verlag, 1980. Translated from the Russian by A. B. Aries.
- [6] N. V. Krylov, "On the rate of convergence of finite-difference approximations for Bellman's equations," *Algebra i Analiz*, vol. 9, no. 3, pp. 245–256, 1997.

-
- [7] N. V. Krylov, "Approximating value functions for controlled degenerate diffusion processes by using piece-wise constant policies," *Electronic Journal of Probability*, vol. 4, no. 2, pp. 1–19, 1999.
 - [8] N. V. Krylov, "On the rate of convergence of finite-difference approximations for Bellman's equations with variable coefficients," *Probab. Theory Related Fields*, vol. 117, no. 1, pp. 1–16, 2000.
 - [9] G. Barles and E. R. Jakobsen, "Error bounds for monotone approximation schemes for Hamilton-Jacobi-Bellman equations," *SIAM J. Numer. Anal.*, vol. 43, no. 2, pp. 540–558 (electronic), 2005.
 - [10] G. Barles and E. R. Jakobsen, "On the convergence rate of approximation schemes for Hamilton-Jacobi-Bellman equations," *M2AN Math. Model. Numer. Anal.*, vol. 36, no. 1, pp. 33–54, 2002.
 - [11] N. V. Krylov, "The rate of convergence of finite-difference approximations for Bellman equations with Lipschitz coefficients," *Appl. Math. Optim.*, vol. 52, no. 3, pp. 365–399, 2005.
 - [12] N. V. Krylov, "A priori estimates of smoothness of solutions to difference Bellman equations with linear and quasilinear operators," *arXiv:math*, vol. 1, no. 1, pp. 1–35, 2006.
 - [13] E. R. Jakobsen, "On the rate of convergence of approximation schemes for Bellman equations associated with optimal stopping time problems," *Math. Models Methods Appl. Sci.*, vol. 13, no. 5, pp. 613–644, 2003.
 - [14] E. R. Jakobsen and K. H. Karlsen, "Convergence rates for semi-discrete splitting approximations for degenerate parabolic equations with source terms," *BIT*, vol. 45, no. 1, pp. 37–67, 2005.
 - [15] E. R. Jakobsen, K. H. Karlsen, and C. La Chioma, "Error estimates for approximate solutions to Bellman equations associated with controlled jump-diffusions," http://www.math.uio.no/eprint/pure_math/2005/pure_2005.html, 2005.
 - [16] I. H. Biswas, E. R. Jakobsen, and K. H. Karlsen, "Error estimates for finite difference-quadrature schemes for a class of nonlocal Bellman equations with variable diffusion," http://www.math.uio.no/eprint/pure_math/2006/pure_2006.html, 2006.
 - [17] I. Gyöngy and D. Šiška, "On randomized stopping," *arXiv:math*, no. 1, pp. 1–11, 2007.
 - [18] H. Dong and N. V. Krylov, "On the rate of convergence of finite-difference approximations for Bellman equations with constant coefficients," *Algebra i Analiz*, vol. 17, no. 2, pp. 108–132, 2005.